



Comparative Morpho-Anatomical Studies of Two Philippine Endemic Species of *Amyema* Tiegh. (Loranthaceae)

Authors:

Romeo M. Tubongbanua, Jr.*, Noe P. Mendez and Victor B. Amoroso

*Correspondence: rmtjr1993@gmail.com

Submitted: 20 January 2022; **Accepted:** 11 August 2022; **Published:** 31 March 2023

To cite this article: Romeo M. Tubongbanua Jr., Noe P. Mendez and Victor B. Amoroso (2023). Comparative morpho-anatomical studies of two Philippine endemic species of *Amyema* Tiegh. (Loranthaceae). *Tropical Life Sciences Research* 34(1): 139–150. <https://doi.org/10.21315/tlsr2023.34.1.9>

To link to this article: <https://doi.org/10.21315/tlsr2023.34.1.9>

Highlights

- The two *Amyema* species differ in the colour of flowers in which pink for *A. curranii* while red for *A. seriata*.
- The two species differ in haustorial attachment: *A. curranii* has several haustoria and creeping, and ovary wall has a presence of hairs, while *A. seriata* has a solitary attachment and the ovary wall that is smooth.
- This is the first morpho-anatomical study of species in genus *Amyema* in the Philippines, which can be used to scrutinize taxa for species delineation.

TLSR, 34(1), 2023

© Penerbit Universiti Sains Malaysia, 2023