

ORIGINAL ARTICLE

Volume 18 No. 2 2026

DOI: 10.21315/eimj2026.18.2.4

ARTICLE INFO

Submitted: 29-03-2025

Accepted: 25-05-2025

Online: 30-06-2026

Translation and Validation of a Chinese Version of the Medical Professionals' Resilience Scale

Shiyue Wang^{1,2}, Garry Kuan¹, Ke Zhou³, Le Li^{1,2}, Mengyuan Zhao¹, Yee Cheng Kueh⁴

¹*Exercise and Sports Science Programme, School of Health Sciences, Universiti Sains Malaysia, Kubang Kerian, Kelantan, MALAYSIA*

²*School of Rehabilitation Medicine, Jiangsu Vocational College of Medicine, Yancheng, Jiangsu, CHINA*

³*Department of Physical Education, Henan University, Kaifeng, Henan, CHINA*

⁴*Biostatistics and Research Methodology Unit, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian, Kelantan, MALAYSIA*

To cite this article: Wang S, Kuan G, Zhou K, Li L, Zhao M, Kueh YC. Translation and validation of a Chinese version of the medical professionals' resilience scale. *Education in Medicine Journal*. 2026;18(2):47–58. <https://doi.org/10.21315/eimj2026.18.2.4>

To link to this article: <https://doi.org/10.21315/eimj2026.18.2.4>

ABSTRACT

Due to its strong psychometric properties, the Medical Professionals Resilience Scale (MeRS) has shown promising results in measuring the resilience of medical professionals. However, a validated Chinese version of the MeRS has yet to be developed. This study aimed to assess the reliability and validity of the Chinese-translated version of the MeRS (MeRS-C) among medical students in Yancheng City, China. The study involved 782 medical student volunteers from Jiangsu Vocational College of Medicine, aged between 19 and 23. The MeRS-C comprises 37 items categorised into four factors and was meticulously translated and adapted into Chinese. An exploratory factor analysis (EFA) was conducted on data from group 1 ($n = 391$) to evaluate the factor structure, and a second-order confirmatory factor analysis (CFA) was conducted on data from group 2 ($n = 391$) to assess model fit. Before being administered to participants, the questionnaire was reviewed by experts and pre-tested. The results of the EFA supported the validity of the MeRS-C, explaining 71.431% of the total variance, with factor loadings ranging from 0.775 to 0.892. The CFA displayed favourable model fit indices (CMICMIN/DF = 1.756; Tucker-Lewis Index (TLI) = 0.96; Standardized Root Mean Square Residual (SRMR) = 0.037; Comparative Fit Index (CFI) = 0.963; and Root Mean Square Error of Approximation (RMSEA) = 0.044) and demonstrated good convergent and discriminant validity. The questionnaire exhibited high internal consistency, with reliability values ranging from 0.897 to 0.967. Thus, the concise Chinese version of the MeRS showed good reliability and validity, making it suitable for identifying sources of resilience in Chinese healthcare trainees.

Keywords: MeRS, Translation, Validation, Resilience

CORRESPONDING AUTHOR

Garry Kuan, Exercise and Sports Science Programme, School of Health Sciences, Universiti Sains Malaysia, 16150 Kubang Kerian, Kelantan, Malaysia

Email: garry@usm.my

INTRODUCTION

Numerous studies have shown that heightened levels of burnout and depression are prevalent among medical students (1–3). Particularly during internships, medical students are confronted with a fresh set of stressors, including the emotional weight of patient deaths, uncertainties in clinical settings and learning processes, interpersonal dynamics with colleagues, and the need to adapt to evolving work and learning environments (4, 5).

Resilience, a construct variously characterised as an intrinsic personality trait or an adaptive process in the face of challenges, emerges during different life stages in response to demanding circumstances (6, 7). The role of resilience in averting and recovering from psychiatric conditions is pivotal (8). Extensive research has consistently highlighted the positive impact that resilience can have on both medical practitioners and their patients (9, 10).

While several scales exist to measure mental resilience, including the Connor–Davidson Resilience Scale (CD-RISC), the Brief Resilience Scale (BRS), the Resilience Scale for Adults (RSA), the Resilience Scale (RS-14), the Post-Traumatic Growth Inventory (PTGI), and the Ego Resiliency Scale (ERS) (11, 12), none of these tools were explicitly designed for healthcare practitioners. The Medical Professionals Resilience Scale (MeRS) was specifically developed to gauge resilience in healthcare providers (13). Comprising 37 items grouped into control, resourcefulness, growth and involvement domains, the MeRS employs a 4-point Likert scale for rating responses (1 = strongly disagree, 2 = disagree, 3 = agree and 4 = strongly agree). A higher MeRS score indicates greater resilience, while a lower score signifies less resilience. The MeRS has been validated among health professionals in Malaysia and Pakistan, with the total number of items ranging from 14 to 37 (13, 14). Based on the confirmatory factor analysis results, these studies have found that this scale has good reliability and validity.

Thus, this study sought to evaluate the reliability and validity of the Chinese version of the MeRS questionnaire (MeRS-C). The study postulated that the MeRS-C questionnaire would demonstrate significant applicability and efficacy, positioning it as a robust instrument for investigating the resilience levels of Chinese medical student trainees.

METHODS

Procedures and Participants

The research was conducted in July 2022 at Jiangsu Vocational College of Medicine in Yancheng City, China, using a cross-sectional study design. Students in their third year of healthcare training across various disciplines such as clinical studies, nursing, rehabilitation therapy, stomatology, and others were invited to participate in the research.

The online survey was conducted via WJX (<https://www.wjx.cn/>), an open-source platform for online surveys that ensures complete respondent anonymity. Upon distribution, the software automatically generates survey access links for all participants. Subsequently, upon survey completion, all identifiable links between participants and their responses are expunged, ensuring that only de-identified data is made available to researchers to safeguard participant anonymity.

Adherence to ethical guidelines was paramount throughout the study, aligning with the ethical standards stipulated in the Declaration of Helsinki (15) and the regulations set forth by Jiangsu Vocational College of Medicine's Ethics Committee to uphold the integrity of the research process and the well-being of study participants.

Measures

Demographic information

During the study, participants were required to provide demographic information, including their age, gender, and field of study. The inclusion criteria were individuals aged 18 to 30 years who voluntarily consented to participate in the research. They were healthcare students from the Jiangsu Vocational College of Medicine. Conversely, participants were excluded from the study if their questionnaire completion time was under 200 seconds or if they provided identical responses to different questions. These criteria were implemented to ensure the reliability and validity of the data collected and to maintain the integrity of the study results.

MeRS

The MeRS, initially developed by Rahman et al. (13) and subsequently validated within the Malaysian medical professional population (13), comprises 37 items organised into four distinct factors: control, resourceful, involvement, and growth. The number of items for each factor is 6, 4, 12, and 15, respectively. Control refers to maintaining composure in challenging situations, influenced by one's inner cognitive processes; resourceful denotes the ability to identify solutions using available means; involvement pertains to effectively managing adversity; while growth signifies personal development amid challenges. Respondents rate these items on a 4-point Likert scale ranging from 1 (strongly disagree) to 4 (strongly agree).

The MeRS demonstrated satisfactory content and face validity while exhibiting a high level of internal consistency. Individual item content validity index (CVI) and face validity index (FVI) values exceeded 0.80. Factor loading values for the final MeRS items ranged from 0.41 to 0.76, with resilience domain Cronbach's alpha coefficients ranging from 0.72 to 0.89. Notably, a higher MeRS score indicates elevated resilience, with lower scores suggesting the opposite. These outcomes underscore the robust psychometric properties of the MeRS, affirming its utility as a reliable and valid instrument for assessing resilience among healthcare professionals.

Cross-cultural translation of the MeRS

The process of forward- and backwards-translating and culturally adapting the original Medical Professionals Resilience Scale (MeRS) into Chinese underwent multiple stages to ensure reliability and validity. As a first step, communication was initiated with the authors of the MeRS to procure the English version of the questionnaire. Following this, a multi-stage approach, outlined by Chai et al. (16) and Yao et al. (17), was employed for translation and cultural adaptation.

Two bilingual and bicultural translators, both native Chinese speakers and well-versed in psychology, translated the MeRS into Chinese individually. Subsequently, the author compared the two forward translations, enabling them to address any discrepancies and ambiguities in collaboration with the translators. This iterative process led to the creation of a preliminary Chinese version of the MeRS.

To validate the translated version, a third bilingual and bicultural translator back-translated the preliminary Chinese questionnaire into English. Notably, the back-translator, a native English speaker without a background in psychology, provided an unbiased perspective. The authors then meticulously compared the back-translated version with the original MeRS to ensure conceptual, semantic, and content equivalence between the two English versions.

In the final validation step, a pilot test was conducted online with ten native Chinese-speaking medical students, representing the target demographic for the Chinese MeRS. These participants were asked to complete the Chinese MeRS without access to the English version. Subsequently, they provided feedback on the clarity of the instructions, the response format, and the overall comprehensibility of the items. The feedback indicated that the Chinese MeRS was clear, easy to comprehend, and swift to complete. Consequently, no further modifications were deemed necessary based on the positive outcomes of the pilot test.

Statistical analysis

Descriptive statistics were utilised to analyse the sample characteristics and item distribution within the study. Construct validity was assessed through both exploratory factor analysis (EFA) and confirmatory factor analysis (CFA). The cohort of 782 participants was divided randomly into two equal groups for data analysis purposes: one group for EFA consisting of 391 participants and another for CFA also comprising 391 participants.

The initial exploratory analysis and assessment of factor reliability were conducted using Statistical Package for the Social Science (SPSS) 28.0 software, while the subsequent CFA was executed utilising Mplus 8.3. Maximum likelihood (ML) is a common estimation method used by CFA, which assumes that the data is multivariate normal distribution (18). The ML method has good statistical properties such as consistency and unbiasedness when the sample is large and the data satisfy the assumption of normal distribution (19). When the data deviate from the normal distribution, the robust maximum likelihood (MLR) method should be chosen (20). MLR can handle non-normally distributed data and provide more robust parameter estimates (21). Since the data did not conform to a normal distribution, the maximum likelihood with MLR approach was adopted to estimate the parameters of the model. To achieve good psychometric properties, factors with high standardised factor loadings (> 0.4) were selected (22).

Based on the four-factor structure and 37-item measurement model of this study, the fit indices used and the recommended fit values are as follows: Chi-square divided by degrees of freedom (χ^2/df) < 3 , Comparative Fit Index (CFI) > 0.90 , the Tucker-Lewis Index (TLI) > 0.90 , Root Mean Square Error of Approximation (RMSEA) < 0.08 , Standardized Root Mean Square Residual (SRMR) < 0.08 (23).

The Cronbach's alpha coefficient (CA) represents the internal consistency reliability of a scale and is the most commonly used measure in scientific research (24). CA > 0.8 indicates good internal consistency of the scale (25). However, CA tends to underestimate the reliability of the underlying structure (26). The minimum acceptable range for composite reliability (CR) is ≥ 0.60 (23). Therefore, this study will use both CA and CR to assess the internal consistency of the scale. An average variance extracted (AVE) value of 0.50 or higher is considered acceptable (27). Discriminant validity was assessed by examining the correlations between the factors in the model. Discriminant validity is established when the correlations between the factors are less than 0.85 (28).

RESULTS

Demographics

A total of 857 participants initially completed the online survey. Subsequently, after screening for validity, 782 questionnaires were found acceptable for analysis, resulting in a response rate of 91.25%. The average age among the participants stood at 21.3 years, with a standard deviation of 2.0 years. Among the respondents, 365 were male, constituting 46.7% of the sample, while 417 were female, making up the remaining 53.3% (refer to Table 1 for demographic information).

Table 1: Participants' demographic information

Variables	Frequency (n)	%
Gender		
Female	417	53.3
Male	365	46.7
Subject/major		
Clinical medicine	152	19.4
Dentistry	97	12.4
Rehabilitation therapy	223	28.5
Pharmacy	62	7.9
Nursing	248	31.8

Reliability Analysis

The Cronbach's alpha coefficient for the MeRS-C scale, calculated with a sample size of 782 participants, yielded a high value of 0.942. Furthermore, the individual Cronbach's alpha coefficients for the growth, involvement, control, and resourceful factors were notably strong at 0.967, 0.956, 0.948, and 0.897, respectively. These values surpassed the standard threshold of 0.7, signifying robust internal consistency reliability within the variables assessed (see Table 2).

Table 2: Reliability analysis

Domains	Cronbach's alpha	Items
Growth	0.967	15
Involvement	0.956	12
Control	0.948	6
Resourceful	0.897	4
Overall	0.942	37

Exploratory Factor Analysis

Exploratory factor analysis (EFA) was conducted on a dataset of 391 entries, revealing a Kaiser-Meyer-Olkin (KMO) value of 0.952. Significantly, the Bartlett spherical test yielded a statistically significant result ($\chi^2 = 17636.247$, $df = 666$, $p < 0.001$), affirming suitability for factor analysis.

Given the MeRS's focus on measuring four dimensions of resilience, we extracted four common factors to align the Chinese version of the questionnaire with the original. Using principal component analysis for factor extraction, we identified common factors with eigenvalues exceeding 1. Notably, the cumulative variance explained by these four common factors stood at 71.431%.

Using the maximum variance orthogonal rotation, the factor loadings across the questionnaire ranged from 0.775 to 0.892. Specifically, factor loadings for each dimension were as follows: growth (factor 1)=0.792–0.864; involvement (factor 2)=0.775–0.853; control (factor 3)=0.849–0.892; resourceful (factor 4)=0.780–0.867. These findings underscored the robustness and dimensionality of the MeRS in capturing various facets of resilience among healthcare professionals.

Confirmatory factor analysis

To further validate the questionnaire, confirmatory factor analysis (CFA) was conducted using Mplus 8.3, building upon the insights outlined in the EFA and involving a sample of 391 participants. The fit indices derived from this analysis were outlined as follows: CMIN/DF = 1.756, RMSEA = 0.044, CFI = 0.963, TLI = 0.96, and SRMR = 0.037 (refer to Table 3 for detailed results). All of these fit indices met general research standards, indicating a favourable model fit for the questionnaire.

Table 3: Goodness-of-fit indexes of the four-factor model for the Chinese version of MeRS (n = 391)

Model	CMIN	DF	CMIN/DF	SRMR	TLI	CFI	RMSEA
Fitness criteria	–	–	< 3	< 0.08	> 0.9	> 0.9	< 0.08
4-factor model	1093.921	623	1.756	0.037	0.96	0.963	0.044

The validation process of the scale involved assessing its reliability through CR and evaluating its convergent validity through AVE. Notably, the factor loadings of the items corresponding to the four resilience factors – growth, involvement, control, and resourceful hopelessness – surpassed 0.6, signifying a high level of item representativeness within the scale.

Additionally, all 4 dimensions exhibited AVE values ranging from 0.645 to 0.758, exceeding the 0.5 benchmark, and CR values ranging from 0.904 to 0.968, surpassing the 0.7 threshold. These outcomes underscored the strong convergent validity of the MeRS-C, indicating that the scale effectively measures the intended constructs with high reliability and consistency.

Table 4: Convergent validity of the Chinese version of the MeRS (n = 391)

Factor	Item	STD. estimate	S.E.	Est./S.E.	CR	AVE
Growth	Q1	0.812	0.018	44.890	0.968	0.669
	Q2	0.825	0.017	48.384		
	Q3	0.807	0.019	43.485		
	Q4	0.787	0.020	39.093		
	Q5	0.836	0.016	51.883		
	Q6	0.783	0.020	38.352		
	Q7	0.789	0.020	39.582		
	Q8	0.841	0.016	53.549		
	Q9	0.833	0.016	50.844		
	Q10	0.772	0.021	36.220		
	Q11	0.831	0.017	50.218		
	Q12	0.776	0.021	36.871		
	Q13	0.862	0.014	61.947		
	Q14	0.839	0.016	52.974		
	Q15	0.864	0.014	62.717		
Involvement	Q16	0.841	0.016	52.458	0.956	0.645
	Q17	0.816	0.018	44.961		
	Q18	0.869	0.014	63.312		
	Q19	0.819	0.018	45.704		
	Q20	0.744	0.024	31.272		
	Q21	0.836	0.017	50.650		
	Q22	0.830	0.017	48.795		
	Q23	0.794	0.020	39.898		
	Q24	0.749	0.023	31.969		
	Q25	0.762	0.022	33.965		
	Q26	0.784	0.021	37.809		
	Q27	0.779	0.021	36.995		
Control	Q28	0.840	0.016	51.220	0.949	0.758
	Q29	0.859	0.015	58.009		
	Q30	0.863	0.014	59.739		
	Q31	0.898	0.012	77.830		
	Q32	0.904	0.011	82.046		
	Q33	0.857	0.015	57.024		

(Continued on next page)

Table 4: (Continued)

Factor	Item	STD. estimate	S.E.	Est./S.E.	CR	AVE
Resourceful	Q34	0.762	0.023	32.927	0.904	0.703
	Q35	0.879	0.015	60.445		
	Q36	0.738	0.025	29.742		
	Q37	0.956	0.010	93.810		

Note: p -value > 0.001 for all items

Discriminant validity was assessed using the correlational relationships among the various factors. Table 5 presents the correlation coefficients among the model's factors. All the correlations are below the recommended value of 0.85, indicating that the four factors possess good discriminant validity.

Table 5: Correlations between latent variables for MeRS

Factor	Growth	Involvement	Control	Resourceful
Growth	1	–	–	–
Involvement	0.467	1	–	–
Control	0.626	0.569	1	–
Resourceful	0.482	0.618	0.617	1

DISCUSSION

The medical profession places significant demands on medical students due to the specialised nature of the field, resulting in the widespread experience of substantial stress and occupational burnout among them (29). These pressures can lead to various negative effects, including reduced empathy, diminished professional motivation, fatigue, deteriorating health, substance abuse, psychological distress and increased suicide rates (30–32). In this context, resilience is crucial, as it helps medical students manage stressors. Individuals with robust psychological resilience can better navigate challenges and stressors in their professional lives, including long work hours, significant responsibilities and emotional strain (33, 34). As future healthcare providers, the psychological resilience of medical students has a direct impact on their performance and adaptability in healthcare environments. Therefore, evaluating the psychological resilience of medical students is essential for their personal growth and for advancing the healthcare sector. Strengthening and nurturing the psychological resilience of medical students enhances not only their ability to manage work pressures and challenges but also the quality and efficiency of healthcare services, which has a significant impact on societal well-being (35). The scale's original authors meticulously analysed the psychological measurement properties of 21 resilience scales. They revamped the resilience concept by amalgamating three resilience perspectives (traits, processes and outcomes) into a novel conceptual framework, culminating in a holistic resilience model featuring four key themes such as growth, control, involvement and resourcefulness. This model conceptualises the relationship between resilience and present and future states, as well as between internal and external conditions, as an inseparable continuum.

The current study is the first to translate the MeRS into Chinese and test its reliability and validity with a cohort of college medical students. The findings revealed that the Chinese MeRS demonstrated strong internal consistency, reliability, construct validity and convergent validity. The study investigated the scale's reliability by examining internal consistency, which resulted in a Cronbach's alpha coefficient of 0.942 for the overall scale and coefficients ranging between 0.897 and 0.967 for the individual dimensions. Comparisons with the CD-RISC (Cronbach's alpha of 0.794), the BRS (Cronbach's alpha of 0.80) and the ARS (Cronbach's alpha of 0.82), indicate reliability comparable with that of other resilience measurement scales (36–38).

Regarding construct validity, the study utilised both EFA and CFA techniques. Bartlett's Spherical Test exhibited a statistically significant result ($\chi^2 = 17636.247$, $p < 0.001$), with a KMO value exceeding 0.5, indicating the appropriateness of factor analysis. Principal component analysis and orthogonal rotation were conducted, elucidating four common factors that explained 71.431% of the total variance. The model fit indices, including CMICMIN/DF, TLI, CFI, and RMSEA, all met the standard thresholds, indicating a robust fit. The factor loadings of the items in the four sub-scales were generally consistent with the analysis conducted by the original authors, indicating acceptable construct validity (13). In a study conducted in Pakistan, the fit indices for the MeRS-37 were relatively poor, possibly due to differences in demographic characteristics, healthcare personnel, and medical students' work environments, all of which could have affected the results (14).

The current study assessed convergent validity by computing the AVE and CR, and evaluated discriminant validity using the square root of AVE. The results of the discriminant validity tests demonstrated that each factor of the MeRS was distinct and did not greatly overlap with the other factors. The research outcomes validated the questionnaire's structural validity and discriminant validity, emphasising the efficiency of the Chinese MeRS as a tool for evaluating the resilience of Chinese medical professionals and healthcare trainees.

Despite the valuable insights gained from this study, some limitations are noted. The reliance on convenience sampling of college students in Jiangsu Province may restrict the generalisability of the findings to a broader population of Chinese medical students. Furthermore, the lack of matching on sociodemographic characteristics, such as gender, could introduce biases. To enhance the robustness and inclusivity of the results, future research should strive to increase sample diversity and demographic balance.

CONCLUSION

The Chinese version of the MeRS was found to be a reliable and valid instrument suitable for assessing resilience factors among Chinese healthcare trainees. Its robust psychometric properties lend credibility to its utility as an effective tool for assessing the resilience levels of Chinese healthcare professionals.

ACKNOWLEDGEMENTS

We gratefully acknowledge all the participants for their cooperation.

ETHICAL APPROVAL

In addition, the study also received ethical approval from the Universiti Sains Malaysia Human Research Ethics Committee (USM/JePEM/22050283).

REFERENCES

1. Dyrbye LN, Massie Jr FS, Eacker A, Harper W, Power D, Durning SJ, et al. Relationship between burnout and professional conduct and attitudes among US medical students. *JAMA*. 2010;304(11):1173–80. <https://doi.org/10.1001/jama.2010.1318>
2. Kuan G, Abdullah N, Kueh YC, Ibrahim MI, Shafei MN. Co-curricular activities and motives for participating in physical activity among health science students at Universiti Sains Malaysia. *Malays J Med Sci*. 2019;26(1):138–46. <https://doi.org/10.21315/mjms2019.26.1.13>
3. Rotenstein LS, Ramos MA, Torre M, Segal JB, Peluso MJ, Guille C, et al. Prevalence of depression, depressive symptoms, and suicidal ideation among medical students: a systematic review and meta-analysis. *JAMA*. 2016;316(21):2214–36. <https://doi.org/10.1001/jama.2016.17324>
4. Dyrbye LN, Thomas MR, Shanafelt TD. Medical student distress: causes, consequences, and proposed solutions. *Mayo Clin Proc*. 2005;80(12):1613–22. <https://doi.org/10.4065/80.12.1613>
5. Haglund MEM, aan het Rot M, Cooper NS, Nestadt PS, Muller D, Southwick SM, et al. Resilience in the third year of medical school: a prospective study of the associations between stressful events occurring during clinical rotations and student well-being. *Acad Med*. 2009;84(2):258–68. <https://doi.org/10.1097/ACM.0b013e31819381b1>
6. Ting LK, Kuan G, Ariffin WN, Cheng KY. Psychometric properties of the self-efficacy scale among undergraduate students in Malaysia. *Malays J Med Sci*. 2019;26(3):119–28. <https://doi.org/10.21315/mjms2019.26.3.10>
7. Taku K. Relationships among perceived psychological growth, resilience and burnout in physicians. *Pers Individ Dif*. 2014;59:120–3. <https://doi.org/10.1016/j.paid.2013.11.003>
8. Liew GC, Kuan G, Chin N-S, Hashim H. Mental toughness in sport: systematic review and future. *Ger J Exerc Sport Res*. 2019;1–14. <https://doi.org/10.1007/s12662-019-00603-3>
9. Mills J, McKimm J. Resilience: why it matters and how doctors can improve it. *Br J Hosp Med*. 2016;77(11):630–3. <https://doi.org/10.12968/hmed.2016.77.11.630>
10. Profit J, Sharek PJ, Amspoker AB, Kowalkowski MA, Nisbet CC, Thomas EJ, et al. Burnout in the NICU setting and its relation to safety culture. *BMJ Qual Saf*. 2014;23(10):806–13. <https://doi.org/10.1136/bmjqs-2014-002831>
11. Runyan C, Savageau JA, Potts S, Weinreb L. Impact of a family medicine resident wellness curriculum: a feasibility study. *Med Educ Online*. 2016;21(1):30648. <https://doi.org/10.3402/meo.v21.30648>

12. Rotenstein LS, Torre M, Ramos MA, Rosales RC, Guille C, Sen S, et al. Prevalence of burnout among physicians: a systematic review. *JAMA*. 2018;320(11):1131–50. <https://doi.org/10.1001/jama.2018.12777>
13. Rahman MA, Yusoff MSB, Roslan NS, Mohammad JA-M, Ahmad A. Development and validation of the medical professionals resilience scale. *BMC Health Serv Res*. 2021;21:482. <https://doi.org/10.1186/s12913-021-06542-w>
14. Sohail M, Yusoff MSB, Hadie SNHB, Othman AB, Rauf H. Psychometric properties of Medical Professionals Resilience Scale (MERS) for medical doctors in Pakistan. *Res Square*. 2022;1–13. <https://doi.org/10.21203/rs.3.rs-1329420/v1>
15. World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*. 2013;310(20):2191–4. <https://doi.org/10.1001/jama.2013.281053>
16. Chai S, Kueh YC, Yaacob NM, Kuan G. Psychometric properties of the Malay version goal content for exercise questionnaire among undergraduate students in health campus, Universiti Sains Malaysia. *Malays J Med Sci*. 2019;26(1):115–24. <https://doi.org/10.21315/mjms2019.26.1.11>
17. Yao L, Zhou K, Zhou Y, Kueh YC, Liu H, Liu Z, et al. Psychometric properties of the Chinese version of the exercise self-efficacy scale for the transtheoretical model: a confirmatory analysis among Chinese children and adolescents. *BMC Public Health*. 2024;24(1):1–11. <https://doi.org/10.1186/s12889-023-17596-2>
18. Gao C, Shi D, Maydeu-Olivares A. Estimating the maximum likelihood root mean square error of approximation (RMSEA) with non-normal data: a Monte-Carlo study. *Struct Equ Model: Multidiscip J*. 2020;27(2):192–201. <https://doi.org/10.1080/10705511.2019.1637741>
19. Kline RB. Principles and practice of structural equation modeling, 4th ed. New York, NY: Guilford Press; 2016.
20. Zhong X, Yuan KH. Bias and efficiency in structural equation modelling: maximum likelihood versus robust methods. *Multivariate Behav Res*. 2011;46(2):229–65. <https://doi.org/10.1080/00273171.2011.558736>
21. Shi D, DiStefano C, Zheng X, Liu R, Jiang Z. Fitting latent growth models with small sample sizes and non-normal missing data. *Int J Behav Dev*. 2021;45(2):179–92. <https://doi.org/10.1177/0165025420979365>
22. DeVon HA, Block ME, Moyle-Wright P, Ernst DM, Hayden SJ, Lazzara DJ, et al. A psychometric toolbox for testing validity and reliability. *J Nurs Scholarsh*. 2007;39(2):155–64. <https://doi.org/10.1111/j.1547-5069.2007.00161.x>
23. Hair JF, Black WC, Babin BJ, Anderson RE. Multivariate data analysis. Boston, MA: Cengage; 2019.
24. Wang J, Wang X. Structural equation modeling: applications using Mplus. Hoboken, NJ: John Wiley & Sons; 2019.
25. Lance CE, Butts MM, Michels LC. The sources of four commonly reported cutoff criteria: what did they really say? *Organ Res Methods*. 2006;9(2):202–20. <https://doi.org/10.1177/1094428105284919>
26. Falk CF, Savalei V. The relationship between unstandardized and standardized alpha, true reliability, and the underlying measurement model. *J Pers Assess*. 2011;93(5):445–53. <https://doi.org/10.1080/00223891.2011.594129>

27. Fornell C, Larcker DF. Evaluating structural equation models with unobservable variables and measurement error. *J Mark Res.* 1981;18(1):39–50. <https://doi.org/10.2307/3151312>
28. Brown TA. *Confirmatory factor analysis for applied research.* New York, NY: Guilford Press; 2006.
29. Ishak W, Nikraves R, Lederer S, Perry R, Ogunyemi D, Bernstein C. Burnout in medical students: a systematic review. *Clin Teach.* 2013;10(4):242–5. <https://doi.org/10.1111/tct.12014>.
30. Ishak WW, Lederer S, Mandili C, Nikraves R, Seligman L, Vasa M, et al. Burnout during residency training: a literature review. *J Grad Med Educ.* 2009;1(2):236–42. <https://doi.org/10.4300/JGME-D-09-00054.1>
31. Benbassat J, Bauml R, Chan S, Nirel N. Sources of distress during medical training and clinical practice: suggestions for reducing their impact. *Med Teach.* 2011;33(6):486–90. <https://doi.org/10.3109/0142159X.2010.531156>
32. Hojat M, Vergare MJ, Maxwell K, Brainard G, Herrine SK, Isenberg GA, et al. The devil is in the third year: a longitudinal study of erosion of empathy in medical school. *Acad Med.* 2009;84(9):1182–91. <https://doi.org/10.1097/ACM.0b013e3181b17e55>
33. Farquhar J, Kamei R, Vidyarthi A. Strategies for enhancing medical student resilience: student and faculty member perspectives. *Int J Med Educ.* 2018;9:1–6. <https://doi.org/10.5116/ijme.5a46.1ccc>
34. Thompson G, McBride RB, Hosford CC, Halaas G. Resilience among medical students: the role of coping style and social support. *Teach Learn Med.* 2016;28(2):174–82. <https://doi.org/10.1080/10401334.2016.1146611>
35. Bacchi, S., Licinio, J. Resilience and psychological distress in psychology and medical students. *Acad Psychiatry.* 2017;41(2):185–8. <https://doi.org/10.1007/s40596-016-0488-0>
36. Extremera MO, Moreno EO, González MC, Ortega FZ, Ruz RP. Validation of resilience scale (CD-RISC) in elite athletes through a structural equation model. *Retos.* 2017;32:96–100. <https://doi.org/10.47197/retos.v0i32.49910>
37. Furstova J, Kascakova N, Solcova IP, Hasto J, Tavel P. How Czecho-Slovakia bounces back: population-based validation of the brief resilience scale in two central European countries. *Psychol Rep.* 2022;125(5):2807–27. <https://doi.org/10.1177/00332941211029619>
38. Trigueros R, Magaz-González AM, García-Tascón M, Alias A, Aguilar-Parra JM. Validation and adaptation of the academic-resilience scale in the Spanish context. *Int J Environ Res Public Health.* 2020;17(11):3779. <https://doi.org/10.3390/ijerph17113779>