



The Impact of Insecticides on Mycelial Growth of *Metarhizium* spp. and Their Efficacy in Controlling Larvae and Pupae of the House Fly (*Musca domestica* L.)

Authors:

Duanpen Wongsorn, Benya Saenmahayak, Nittaya Pitiwittayakul and Surachai Rattanasuk*

***Correspondence:** surachai_med@hotmail.com

Submitted: 30 April 2024; **Accepted:** 27 January 2025; **Early view:** 7 March 2025

To cite this article: Duanpen Wongsorn, Benya Saenmahayak, Nittaya Pitiwittayakul and Surachai Rattanasuk. (in press). The impact of insecticides on mycelial growth of *Metarhizium* spp. and their efficacy in controlling larvae and pupae of the house fly (*Musca domestica* L.). *Tropical Life Sciences Research*.

Highlights

- The NMMet_SS9/2 and NMMet_CLPK4/1 isolates exhibited the highest infection rates for larvae and pupae, respectively.
- Cypas® 250 EC was less toxic to the hyphal growth of the NMMet_SS9/2 and NMMet_CLPK4/1 isolates than Tanidil®-T.
- The NMMet_SS9/2 and NMMet_CLPK4/1 isolates cultured on PDA with Cypas® 250 EC demonstrated house fly control effectiveness comparable to those grown on PDA alone.

EARLY VIEW

The Impact of Insecticides on Mycelial Growth of *Metarhizium* spp. and Their Efficacy in Controlling Larvae and Pupae of the House Fly (*Musca domestica* L.)

¹Duanpen Wongsorn, ²Benya Saenmahayak, ¹Nittaya Pitiwittayakul and ³Surachai Rattanasuk

¹Department of Plant Science, Faculty of Agricultural Innovation and Technology, Rajamangala University of Technology Isan, Nakhon Ratchasima, 30000 Thailand

²Department of Animal Science, Faculty of Agricultural Innovation and Technology, Rajamangala University of Technology Isan, Nakhon Ratchasima, 30000 Thailand

³Major of Biology, Department of Science and Technology, Faculty of Liberal Arts and Science, Roi Et Rajabhat University, Roi Et 45120 Thailand

Corresponding author: surachai_med@hotmail.com

Running head: The Control of the House Fly Larvae and Pupae

Submitted: 30 April 2024; **Accepted:** 27 January 2025; **Early view:** 7 March 2025

To cite this article: Duanpen Wongsorn, Benya Saenmahayak, Nittaya Pitiwittayakul and Surachai Rattanasuk. (in press). The impact of insecticides on mycelial growth of *Metarhizium* spp. and their efficacy in controlling larvae and pupae of the house fly (*Musca domestica* L.). *Tropical Life Sciences Research*.

Abstract. *Metarhizium* spp. are entomopathogenic hyphomycete fungi with great potential as biological control agents against insects and as a component within integrated pest management systems. This study evaluated 10 *Metarhizium* spp. isolates for their effectiveness against house fly larvae and pupae. The isolates with the highest infection rates were tested for compatibility with insecticides. NMMet_SS9/2 and NMMet_CLPK4/1 were the most effective, with infection rates of 86.67% and 60.00% for larvae and pupae, respectively. Both isolates were cultured on Potato Dextrose Agar (PDA) mixed with Cypas® 250 EC (100 and 50 ml per 6.25 liters) and Tanidil®-T (100 g and 50 g per liter) to assess the impact of insecticides on mycelial growth. After 28 days, the radial growth of NMMet_SS9/2 (81.33 mm) and NMMet_CLPK4/1 (77.67 mm) on PDA with Cypas® 250 EC (50 ml per 6.25 liters) showed no significant differences ($p>0.05$) compared to growth on PDA alone. A spore suspension

(1×10^8 spores/ml) of NMMet_SS9/2 and NMMet_CLPK4/1 cultured on PDA with Cypas® 250 EC (50 ml per 6.25 liters) was used to control house fly larvae and pupae, resulting in mortality rates of 91.11% (NMMet_SS9/2) and 75.56% (NMMet_CLPK4/1), with infection rates of 77.78% and 71.11%, respectively. No significant differences ($p > 0.05$) were observed in mortality or infection rates compared to spores cultured on PDA without insecticides. These findings confirm that NMMet_SS9/2 and NMMet_CLPK4/1 are highly effective against house fly larvae and pupae and can proliferate on media containing Cypas® 250 EC at 50 ml per 6.25 liters without compromising their insecticidal properties, making them promising candidates for integrated pest management strategies.

Keywords: Entomopathogenic Fungi, Biological Control, Pest Management, Pyrethroids

INTRODUCTION

The house fly, scientifically known as *Musca domestica* L. (Diptera: Muscidae), is a pest that carries a multitude of pathogens, which affect both humans and animals worldwide. Houseflies serve as vectors for the mechanical transmission of a wide range of pathogens, including helminthic eggs, protozoan cysts and trophozoites, bacteria, fungi, and viruses, which they disseminate through their vomit or excreta. These insects are implicated in the spread of over 100 types of pathogens, such as those responsible for cholera, anthrax, Shigella infections, ORF virus, and ascariasis (Olagunju 2022). It is typically found in residential areas, waste disposal sites, and locations with decaying food or animal remains (Abbas *et al.* 2013).

House flies can produce more than two broods or generations per year (multivoltine insects). They undergo 10–12 life cycles annually in temperate climates, with their populations peaking during the summer months. These flies neither migrate seasonally, enter diapause, nor survive under adverse environmental conditions. Consequently, their populations align periods of active development and reproduction with seasons when resources are most abundant (Reynolds 2017). Furthermore, the house fly exhibits a high rate of reproduction. It lays approximately 120-150 eggs per batch and can lay eggs 5-20 times, resulting in an average of 500 eggs over its lifetime (Iqbal *et al.* 2014). This rapid reproduction contributes to its status as a significant pest.

Chemical agents are frequently utilized to address house fly infestations due to their convenience and swift control capabilities. Organophosphates, carbamates, and organochlorines are selected for their immediate effectiveness. Organophosphate insecticides, which are esters of phosphoric acid or its sulfur analogs, are extensively utilized for controlling insect pests due to their potent toxicity, rapid environmental degradation, and

selective biological activity. Their mechanism of action involves inhibiting acetylcholinesterase (AChE), an enzyme responsible for breaking down acetylcholine, thereby disrupting neural transmission, inducing hyperexcitation, and ultimately resulting in the death of the target organisms (Barata *et al.* 2004). Pyrethroid insecticides are the most commonly utilized class for controlling house flies due to their high efficacy, low mammalian toxicity, brief environmental persistence, and limited impact on non-target organisms (Elliott 1980).

However, these chemicals can have detrimental effects on users and may linger in produce and the environment. Moreover, insects can develop resistance to these chemicals (Aktar *et al.* 2009; Abbasi *et al.* 2023). Acevedo *et al.* (2009) observed that house flies developed resistance against two commercial insecticides, namely 2,2-dichlorovinyl dimethyl phosphate (DDVP), and permethrin. Additionally, house flies also developed resistance to permethrin, deltamethrin, beta-cypermethrin, and propoxur (Wang *et al.* 2019).

To achieve effective insect control, alternatives such as botanical extracts and various bacterial strains are being investigated as potential replacements or supplements to chemical insecticides. This is in line with the Integrated Pest Management (IPM) approach (Miana *et al.* 2018). Significantly, entomopathogenic fungi, especially *Metarhizium* spp., are viewed as a promising solution for integrated pest management. This is due to their limited host range, safety, environmental compatibility, and the feasibility of mass production. These fungi can target a wide range of insect orders, including Lepidoptera, Hymenoptera, Hemiptera, Homoptera, Heteroptera, Diptera, and Coleoptera (Balachander *et al.* 2009; Saranraj & Arul 2017).

However, challenges emerge when insecticides are combined with entomopathogenic fungi or when fungi are used in areas previously treated with insecticides. Certain chemicals can adversely affect the growth, vitality, and efficiency of fungi. For instance, specific chemical insecticides, such as Profenofos (50 EC), Indoxacarb (14.5% EC), and Methyldemeton, significantly impact fungal growth, while others, like Chlorpyrifos 20 EC, have a milder effect (Amutha *et al.* 2010). Moreover, the use of insecticides at rates divergent from the recommended label rates can also influence fungi (Abidin *et al.* 2017; Pelizza *et al.* 2017). Consequently, it is crucial to study the impact of commonly used insecticides on the efficacy of entomopathogenic fungi in controlling house flies for the development of effective Integrated Pest Management strategies in the future.

This research aimed to study the effects of insecticide on the growth of mycelium of *Metarhizium* spp. and to test the efficacy of *Metarhizium* spp. cultured on media mixed with insecticide in controlling larvae and pupae of the house fly.

MATERIALS AND METHODS

The Effective Screening of *Metarhizium* spp. for Controlling House Flies

The fungal spore preparation

The 10 *Metarhizium* spp. isolates (NMMet_DKT9/1, NMMet_SN2/1, NMMet_CLPK4/1, NMMet_NBM10/1, NMMet_LTMC7/2, NMMet_KTLS7/3, NMMet_BAL7/1, NMMet_NS7/2, NMMet_SS9/2 and NMMet_SS10/3) obtained from plant protection laboratory, Department of Plant Science, RMUTI Thailand were cultured on Potato Dextrose Agar (PDA). The culture was maintained in darkness at a temperature of 28-30°C for 21 days. Subsequently, the spores were harvested using a surfactant (Tween 20®, 0.05%). The fungal culture was diluted in a petri dish, and a sterilized loop was employed to scrape the mycelium and spores, which were then filtered through sterile, double-layered muslin cloth to remove solid debris. The spore suspension was then quantified using a Hemocytometer under a microscope at 40X magnification. The concentration of the spore suspension was adjusted to a density of 1×10^8 spores/ml. The fungal spore preparation method was modified from Bharathi *et al.* (2022).

Insect culture

Adult house flies were sourced from a poultry farm located at the training center of Nong Rawieng, RMUTI, in Nakhon Ratchasima, Thailand (latitude 14°57'42.5"N and longitude 102°10'19.6"E). Sampling methods included net collection and baited traps, and the collection was conducted between 09:00 and 12:00. The collected house flies were subsequently classified following the criteria outlined by Geden *et al.* (2021). They were then reared in mesh cages with dimensions of 30x30x30 cm. The adult house flies were nourished with a synthetic diet comprising powdered milk (25 g) and dry yeast (25 g) and sugar (50 g). Upon laying eggs, the resulting larvae were fed a semi-artificial diet consisting of 50 g of coarse bran, 10 g of fine bran, and 1 g of pineapple fruit. These ingredients were thoroughly mixed using a blender (Ali *et al.* 2024). The rearing process was conducted in a laboratory setting, maintaining a temperature of 25-30°C, following the methodologies prescribed by Mishra *et al.* (2011). Once the larvae reached the third instar (30-32 hours after hatching), one-day-old pupae (24 hours after pupation) were prepared for subsequent testing.

House fly assay

The third instar larvae or one-day-old pupae of the house fly were placed in 20-ounce plastic cups containing a semi-artificial diet mixed with suspended spores of *Metarhizium* spp., with each isolate at a final concentration of 1×10^8 spores/ml. The treatments were compared with two controls: a semi-artificial diet without any additives and a semi-artificial diet containing Tween 20® (0.05%). The experiment was conducted using a Completely Randomized Design (CRD) with three replicates for each treatment. Each replicate (plastic cup) contained 30 house fly larvae or pupae.

After inoculating the *Metarhizium* spp. isolates on larvae or pupae, they were transferred to an incubator in the dark at a temperature of 28-30°C and 60-80% relative humidity (RH.). When dead larvae and pupae were found, the insects' skin surface was treated with a 0.5% Sodium Hypochlorite solution for 3-5 minutes, followed by rinsing with sterile water for 3-5 minutes, repeated twice. The larvae and pupae were then transferred to a moist chamber and incubated in the dark at a temperature of 28-30°C and 60-80% R.H. Daily observations of mortality and fungal infection were recorded for 14 consecutive days. Correction for mortality in the control treatment were done using Abbott's formula (Abbott 1925) as following;

$$[(\text{Test mortality} - \text{Control mortality}) \times 100] / [100 - \text{Control mortality}]$$

The Mycelial Growth of *Metarhizium* spp. on the Culture Medium Mixed with Insecticides

Culture media preparation

The culture medium for the fungal culture was prepared using Potato Dextrose Agar (PDA). Initially, 200 g of potato were boiled in water for 15-20 minutes until the boiling point was reached. The mixture was subsequently filtered to retain only the water. Various additives were then incorporated into the mixture, including 20 g of glucose and 20 g of agar. After autoclaving (at 121°C for 15 minutes) and cooling down the media, the insecticides (Cypas® 250 EC. and Tanidil®-T, as shown in Table 1) were added at two concentrations; the recommended application rate and half the recommended rate. For Cypas® 250 EC, the concentrations were 100 ml and 50 ml per 6.25 liters, and for Tanidil®-T, they were 100 g and 50 g per liter. The mixture was then shaken for 2 minutes to achieve a homogeneous

distribution of the added compounds. This method was modified from Schumacher & Poehling (2012).

Table 1. Chemical insecticides, active ingredients, and recommended application rates.

Commercial name	Ingredient	Chemical class	Recommended application rate	Manufacturer
Cypas® 250 EC	Cypermethrin 25%	pyrethroids	100 mL/6.25 L	Q Fac Company Limited
Tanidil®-T	Coumaphos 3%, Propoxur 2%	organophosphate	100 g/1 L	Bayer

Fungal culture

The NMMet_SS9/2 and NMMet_CLPK4/1, which have been reported to be highly effective in controlling house fly larvae and pupae, respectively, were cultured on PDA for 14 days. Following this, a cork borer with a diameter of 0.7 mm was used to inoculate at the periphery of the colonized area, and they were then transferred to the center of a Petri dish containing the medium. The experiment included five treatments: PDA alone, PDA + Cypas® 250 EC at the recommended application rate (100 ml per 6.25 liters), PDA + Cypas® 250 EC at half the recommended application rate (50 ml per 6.25 liters), PDA + Tanidil®-T at the recommended application rate (100 g per liter), and PDA + Tanidil®-T at half the recommended application rate (50 g per liter). The experimental design followed a CRD, with each treatment replicated five times, and each replicate comprising five culture media plates.

The fungal culture was incubated in the dark at a temperature of 28-30°C and 70-80% RH. The growth of fungus was measured by the colony diameter at intervals of 7, 14, 21, and 28 days post-inoculation. The inhibition of mycelium growth (I) was computed using the following formula:

$$I (\%) = [(Dc-Dt)/Dc] \times 100$$

I (%): inhibition percentage; Dc: average diameter of the control colonies; Dt: average diameter of the treated colonies.

The Effectiveness of NMMet_SS9/2 and NMMet_CLPK4/1 Cultured On Media Containing An Insecticide in Controlling House Fly Larvae and Pupae

The 1x10⁸ spores/ml spore suspension of *Metarhizium* spp. isolates (NMMet_SS9/2 and NMMet_CLPK4/1) cultured on PDA and PDA mixed with Cypas® 250 EC at half the

recommended application rate were mixed with a semi-artificial diet in plastic cups containing house fly larvae or pupae. After inoculating the *M. anisopliae* isolates on larvae or pupae, they were transferred to an incubator in the dark at a temperature of 28-30°C and 60-80% RH. Daily observations of mortality and fungal infection were recorded for 14 consecutive days. Correction for mortality in the control treatment was done using Abbott's formula (Abbott 1925). This experiment was compared with the fungus cultured on PDA, a control method without any insecticides, and Tween20® (0.05%). The experiment was conducted using a CRD with three replicates for each method. Each replicate used 30 house fly larvae or pupae.

DATA ANALYSIS

Statistical analysis was performed, which included an analysis of variance and a comparison of the mean differences among each treatment. This was conducted using Duncan's New Multiple Range Test (DMRT) in conjunction with the Statistical Analysis System (SAS) software Version 9.00. (SAS Institute Inc. 2006)

RESULTS

The 10 isolates of *Metarhizium* spp. were effective in controlling both larvae and pupae of house flies. For larvae control, NMMet_SS9/2 exhibited the highest infection percentage at 86.67%, but there was no statistical difference ($p > 0.05$) in the infection rates of NMMet_BAL7/1 (71.11%) and NMMet_NS7/2 (68.89%) isolates. Moreover, in the pupal test, NMMet_CLPK4/1 isolate showed the highest infection rate at 60.00%, which was statistically different ($p < 0.05$) from NMMet_LTMC7/2 isolate, which had the lowest infection rate at 15.56%. (Fig. 1).




Figure 1. The effectiveness of *Metarhizium* spp. (1×10^8 spores/mL) against 3rd instar larvae and pupae of the house fly (*Musca domestica* L.) was evaluated under laboratory conditions (28–30°C and 60–80% RH). Experiments were performed in triplicate. Error bars represent $\pm$ SD. Infection percentages followed by the same letter are not significantly different ($p > 0.05$) according to DMRT.

The Mycelial Growth of *Metarhizium* spp. on the Culture Medium Mixed with Insecticides

On the 28th day, (Tables 2 and 3), both isolates, NMMet_SS 9/2 and NMMet_CLPK4/1 demonstrated superior growth on PDA mixed with Cypas® 250 EC compared to PDA mixed with Tanidil®-T. The isolate NMMet_SS9/2 was unable to grow on PDA mixed with Tanidil®-T at the recommended application rate, exhibiting a significant mycelium growth inhibition of up to 93.60%. However, it was able to grow on PDA mixed with Tanidil®-T at half the recommended application rate, with a colony diameter of 20.83 mm and a mycelium growth inhibition of 74.99%. In the case of PDA mixed with Cypas® 250 EC at a concentration of 50 ml per 6.25 liters, the fungal colony diameters were 81.33 mm, showing no statistically significant differences ($p < 0.05$) in colony diameter compared to PDA alone (83.33 mm). At the recommended application rate (100 ml per 6.25 liters) for PDA mixed with Cypas® 250 EC, the fungal colony diameters were 49.00 mm, indicating a mycelium growth inhibition of 41.19%. Similarly, the isolate NMMet_CLPK4/1 cultured on PDA exhibited the largest colony diameter (80.50 mm), which was not significantly different ($p > 0.05$) from PDA mixed with Cypas® 250 EC at half the recommended rate (77.67 mm), but was significantly different ($p < 0.05$) from other treatments. Conversely, when the fungus was cultivated on PDA combined with Tanidil®-T at both the recommended and half-recommended rates, it showed limited growth, with a colony diameter of 18.83 mm.

Table 2. Colony diameter and mycelial growth inhibition of the *Metarhizium* spp. isolate NMMet_SS9/2 on culture medium mixed with insecticides.

Treatments	Parameters ^{1/}							
	Colony diameter (mm ± sd)				Mycelium growth inhibition over control (% ± sd)			
	7 days	14 days	21 days	28 days	7 days	14 days	21 days	28 days
PDA (Control)	28.83 ± 1.04 ^a	46.00 ± 2.60 ^a	70.17 ± 1.89 ^a	83.33 ± 0.76 ^a	0.00 ± 0.00 ^d	-	-	-
PDA+Cypas [®] 250 EC. ^{2/}	20.33 ± 1.53 ^b	30.67 ± 0.76 ^c	42.33 ± 2.84 ^c	49.00 ± 1.32 ^b	29.54 ± 2.75 ^c	33.18 ± 4.30 ^c	39.56 ± 5.75 ^c	41.19 ± 2.13 ^{ab}
PDA+Cypas [®] 250 EC. (½) ^{3/}	17.17 ± 2.02 ^c	35.00 ± 1.32 ^b	65.00 ± 1.73 ^b	81.33 ± 1.04 ^a	40.29 ± 8.72 ^b	23.79 ± 4.30 ^d	7.34 ± 2.73 ^d	2.40 ± 0.60 ^c
PDA+Tanidil [®] -T ^{4/}	5.00 ± 0.00 ^d	5.00 ± 0.00 ^e	5.00 ± 0.00 ^e	5.33 ± 0.29 ^e	82.64 ± 0.62 ^a	89.11 ± 0.60 ^a	92.87 ± 0.20 ^a	93.60 ± 0.39 ^a
PDA+Tanidil [®] -T (½) ^{5/}	5.00 ± 0.00 ^d	12.17 ± 1.89 ^d	15.67 ± 0.58 ^d	20.83 ± 1.61 ^d	82.64 ± 0.62 ^a	73.55 ± 3.93 ^b	77.65 ± 1.29 ^b	74.99 ± 2.12 ^a
<i>P-value</i>	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	0.0104

Notes. ^{1/} Means ± sd within a column followed by the same letter are not significantly different (DMRT, $p > 0.05$), sd = standard deviation; ^{2/} Cypas[®] 250 EC 100 mL/6.25 L (recommended application rate); ^{3/} Cypas[®] 250 EC 50 mL/6.25 L (half-recommended application rate); ^{4/} Tanidil[®]-T 100 mL/1 L (recommended application rate); ^{5/} Tanidil[®]-T 50 mL/1 L (half-recommended application rate)

Table 3. Colony diameter and mycelial growth inhibition of the *Metarhizium* spp. isolate NMMet_CLPK4/1 on culture medium mixed with insecticides.

Treatments	Parameters ^{1/}							
	Colony diameter (mm±sd)				Mycelium growth inhibition over control (%±sd)			
	7 days	14 days	21 days	28 days	7 days	14 days	21 days	28 days
PDA (Control)	28.00 ± 0.05 ^a	46.67 ± 1.89 ^a	65.83 ± 2.47 ^a	80.50 ± 3.61 ^a	0.00 ± 0.00 ^d	-	-	-
PDA+Cypas [®] 250 EC ^{2/}	19.00 ± 0.05 ^c	31.17 ± 0.29 ^c	46.83 ± 0.29 ^c	59.00 ± 1.80 ^b	32.14 ± 1.57 ^b	33.15 ± 2.44 ^c	28.80 ± 2.28 ^c	26.55 ± 5.42 ^c
PDA+Cypas [®] 250 EC(^{1/2}) ^{3/}	21.83 ± 1.04 ^b	37.76 ± 1.15 ^b	56.00 ± 0.50 ^b	77.67 ± 1.04 ^a	21.98 ± 4.86 ^c	19.22 ± 3.24 ^d	14.84 ± 3.92 ^d	3.45 ± 2.36 ^d
PDA+Tanidil [®] -T ^{4/}	5.00 ± 0.00 ^d	5.00 ± 0.00 ^e	6.00 ± 0.00 ^e	8.67 ± 0.76 ^d	82.14 ± 0.32 ^a	89.27 ± 0.44 ^a	90.88 ± 0.35 ^a	89.19 ± 1.46 ^a
PDA+Tanidil [®] -T (^{1/2}) ^{5/}	5.00 ± 0.00 ^d	10.50 ± 0.87 ^d	11.67 ± 2.89 ^d	18.83 ± 1.04 ^c	82.14 ± 0.32 ^a	77.51 ± 1.45 ^b	82.33 ± 3.97 ^b	76.54 ± 2.32 ^b
<i>P-value</i>	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001

Notes. ^{1/} Means ± sd within a column followed by the same letter are not significantly different (DMRT, $p > 0.05$), sd = standard deviation; ^{2/} Cypas[®] 250 EC 100 mL/6.25 L (recommended application rate); ^{3/} Cypas[®] 250 EC 50 mL/6.25 L (half-recommended application rate); ^{4/} Tanidil[®]-T 100 mL/1 L (recommended application rate); ^{5/} Tanidil[®]-T 50 mL/1 L (half-recommended application rate)

The characteristics of the fungal colony at 28 days (Fig. 2) on PDA and PDA mixed with Cypas® 250 EC at half the recommended rate demonstrated that the colony and hyphae could smoothly proliferate on the agar surface. The colony on PDA initially presented as white with a slight green hue, transitioning to a green color with a brownish tint as it matured. Conversely, when cultured on PDA mixed with Cypas® 250 EC the hyphae exhibited slower growth, appearing white, with a dense compaction of hyphae around the colony edges. In the case of PDA mixed with Tanidil®-T, the hyphae failed to grow, and on PDA mixed with Tanidil®-T at half the recommended rate, the fungi produced only a limited number of white hyphae, predominantly clustered at the edge of the medium piece.




Figure 2. The colony characteristics of *Metarhizium* spp. isolates NMMet_SS9/2 and NMMet_CLPK 4/1 on PDA mixed with insecticides—Cypas® 250 EC (100 ml and 50 ml/6.25 liters) and Tanidil®-T (100 g and 50 g/liter)—were observed at 21 days.

The Effectiveness of the *Metarhizium* spp. Cultured on Media Mixed with an Insecticide in Controlling House Fly Larvae and Pupae

The NMMet_SS 9/2 isolate cultured on PDA mixed with Cypas® 250 EC (50 ml per 6.25 liters) demonstrated the highest mortality rate of house fly larvae at 93.33%. This rate was not statistically significantly different ($p>0.05$) from the NMMet_SS 9/2 cultured on PDA alone (91.11%). The infection rates were closely comparable at 84.44% and 77.78% for the fungus cultured on PDA and PDA mixed with Cypas®, respectively. Similarly, the NMMet_CLPK4/1 isolate cultured on PDA mixed with Cypas® 250 EC at half the recommended rate exhibited a

mortality rate of 75.56% for house fly pupae. This rate was comparable to the e NMMet_CLPK4/1 cultured on PDA alone (82.22%). The pupal infection rates were 73.33% and 71.11% for the fungus cultured on PDA and PDA mixed with Cypas® 250 EC at half the recommended rate, respectively (Table 4).

Table 4. The effectiveness of the *Metarhizium* spp., isolates NMMet_SS9/2 and NMMet_CLPK4/1, cultured on media mixed with insecticide, in controlling house fly (*Musca domestica* L.).

Treatments	NMMet_SS 9/2		NMMet_CLPK 4/1	
	Larval mortality (% ± sd)	Larval Infection (% ± sd)	Pupal mortality (% ± sd)	Pupal Infection (% ± sd)
Control (non-spraying)	4.44 ± 3.85 ^b	0.00 ± 0.00 ^b	0.00 ± 0.00 ^b	0.00 ± 0.00 ^b
Tween® 20 0.05%	8.89 ± 3.85 ^b	0.00 ± 0.00 ^b	0.00 ± 0.00 ^b	0.00 ± 0.00 ^b
Fungus cultured on PDA	91.11 ± 3.85 ^a	84.44 ± 10.18 ^a	82.22 ± 7.70 ^a	73.33 ± 11.55 ^a
Fungus cultured on PDA+Cypas® 250 EC ^{1/}	93.33 ± 6.67 ^a	77.78 ± 3.85 ^a	75.56 ± 3.85 ^a	71.11 ± 3.85 ^a
P-value	<0.0001	<0.0001	<0.0001	<0.0001

Notes. Means ± sd within a column followed by the same letter are not significantly different (DMRT, $p>0.05$); sd = standard deviation; ^{1/} Cypas® 250 EC 50 ml/6.25 liters of water (half recommended application rate)

DISCUSSIONS

The results of this study indicated that all 10 isolates of *Metarhizium* spp. could infect both larvae and pupae of the house fly. However, the infection percentage varied among each isolate. Additionally, *Metarhizium* spp. could penetrate and destroy larvae of the house fly more rapidly than pupal stages. After a 2-day inoculation period, fungal hyphae were observed on the bodies of house fly larvae, whereas in the pupal stage, hyphae covered the pupae after 10 days of testing (data not shown in the table). This was consistent with Farooq & Feed (2016), who reported that the virulence of insect destruction by fungi depends on the morphological characteristics such as age, sex, and nutrition of the insect. Similarly, Sharififard *et al.* (2011) reported that the LC₅₀ values of *M. anisopliae* for controlling house flies differ depending on the life stage, with LC₅₀ values ranging from 1.65 to 3x10⁶ conidia per gram for adult stage and 7.3 x 10⁴ to 2.9x10⁶ conidia per mL for the larval stage. Additionally, Mishra *et al.* (2011) found that *M. anisopliae* is more effective in controlling house flies in the adult stage than in the larval stage. The time required for fungal penetration and destruction of insects, from spore contact to spore germination on the insect's body, fungal growth within the insect, and extrusion of hyphae, is approximately 96 hours under favorable environmental conditions (temperature and humidity), but takes longer under unfavorable conditions (Mohammed 2018).

The ability of entomopathogenic fungi to control insects depends on several factors, such as the species and virulence of the entomopathogenic fungi, the target insect, the duration of entomopathogenic fungi contact with the insect, as well as environmental factors including temperature, humidity, sunlight, and rainfall (Bugti *et al.* 2020; Quesada Moraga *et al.* 2024). Some species of entomopathogenic fungi have specific host ranges. The destruction of insects by entomopathogenic fungi may be due to secondary metabolites or toxins produced by the fungi. However, some species of entomopathogenic fungi only invade and compete for essential mineral nutrients inside the insect's body to sustain their own life (Bihal *et al.* 2023).

The compatibility of entomopathogenic fungi with insecticides test results revealed that Cypas® 250 EC, a member of the pyrethroids class, exhibited less toxicity to the hyphal growth of *M.anisopliae* compared to Tanidil®-T. When Cypas® 250 EC was applied at half the recommended rate (50 ml per 6.25 liters of water) on both NMMet_SS9/2 and NMMet_CLPK4/1, the fungal hyphae could grow and produce spores. Conversely, when cultured on PDA mixed with Tanidil®-T, the fungi were unable to grow, with hyphal growth only occurring in the region where the fungus was deposited. This finding aligns with the results of Oliveira *et al.* (2003), who reported that Alpha-Cypermethrin, Thiamethoxam, and Cyfluthrin exhibited lower inhibition to conidia germination at both field recommended and half-field recommended rates. Similarly, Oliveira *et al.* (2015) found that a Cypermethrin-6% wettable powder chemical insecticide, at recommended concentration, double recommended concentration, and half recommended concentration, had no effect on the size of colony diameter but reduced spore viability and conidial production compared to the control (without insecticide). Apoorva and Ramaswamy (2013) conducted compatibility testing of *Metarhizium anisopliae* with three organophosphate compounds (phorate, malathion, and chlorpyrifos) and two pyrethroids (deltamethrin and permethrin), finding that phorate was notably more toxic to *M. anisopliae* than the other pesticides, significantly inhibiting both vegetative growth and sporulation. Similarly, Schumacher and Poehling (2012) examined the *in vitro* effects of various pesticide concentrations, including fipronil, imidacloprid, neemazal, and amitraz, on *M. anisopliae*. Their study revealed that only fipronil, an organophosphate, exhibited moderate toxicity to *M. anisopliae* at a concentration of 200 ppm.

The compatibility between insecticides and entomopathogens depends on various factors. For instance, Tamai *et al.* (2002) reported that insecticide products with similar modes of action, produced by different companies, may elicit different responses from pathogens due to variations in the ingredients (inert ingredients and adjuvants) used in each product's formulation. Such variations are contingent on the presence of compounds that block conidia metabolic functions and the concentrations of active compounds (Antonio *et al.* 2001; Kumar *et al.* 2000).

The mechanism of action of insecticides on entomopathogens is described by Oliveira *et al.* (2003). They reported that molecules analogous to prosthetic groups diffuse to the cytoplasm, where they bind to specific receptors, affecting membrane permeability and enzymatic synthesis. Consequently, this process influences metabolic processes, and the same inhibitory mechanism is likely responsible for differences in conidial germination and vegetative growth in *M. anisopliae*. Additionally, the interaction depends on the type and isolate of the fungus, as different isolates of entomopathogenic fungi exhibit varying levels of tolerance to insecticides. Furthermore, these fungi can also degrade the pyrethroid cypermethrin (Oliveira *et al.* 2016).

NMMet_SS9/2 and NMMet_CLPK4/1 cultured on PDA mixed with Cypas® 250 EC at half the recommended rate were used in controlling both larvae and pupae of house flies. Culturing on PDA mixed with Cypas® 250 EC did not affect the efficacy of insect pest control. These findings align with the results of Jantanapim *et al.* (2015), who reported that *B. bassiana* cultured on PDA combined with Thiamethoxam resulted in the highest mortality of cassava mealybug, with no statistical significance ($p>0.05$) compared to the fungus cultured on PDA. Similarly, Pong-Anan *et al.* (2015) reported that *B. bassiana* cultured on PDA combined with Buprofezin caused the highest mortality of brown planthoppers, with no significant difference compared to the fungus cultured on PDA. This may be attributed to the ability of entomopathogenic fungi to degrade insecticides. Oliveira *et al.* (2016) reported that the fungus *B. bassiana* can degrade Cypermethrin. In a similar vein, Ong *et al.* (2019) found that *M. anisopliae* can degrade both insecticides (Chlorpyrifos and Cypermethrin) by more than 80%, which is significantly higher than the control soil (47-61%). Abd El-Ghany and Masmali (2016) demonstrated that *M. anisopliae* can degrade Organophosphorus by more than 90%, sharing the same category and action mode as chlorpyrifos.

In addition to using entomopathogenic fungi cultured on media containing insecticides for pest control, there is also a trend of using these Entomopathogenic fungi in combination with insecticides to control various insect species. The effectiveness of insect control does not decrease when fungi are combined with insecticides. Sharififard *et al.* (2011) combined *M. anisopliae* with Spinosad to control house flies, resulting in a synergistic effect that increased house fly mortality and reduced the lethal time. Similarly, Ong *et al.* (2017) found that the combination of *M. anisopliae* and a mixture of Chlorpyrifos and Cypermethrin (ChCy) resulted in 62-72% house fly larva mortality. Furthermore, Farooq & Freed (2016) used *B. bassiana* in combination with insecticides to control house flies. They discovered that when *B. bassiana* was combined with acetamiprid, Emamectin, Imidacloprid, or lufenuron, there was a significant decrease in longevity, fecundity, egg hatching, percent pupation, pupal weight, and adult emergence. Additionally, larval duration and pupal period were prolonged.

CONCLUSIONS

The NMMet_SS9/2 and NMMet_CLPK4/1 isolates were identified as the most effective in controlling house fly larvae and pupae, respectively. Both isolates exhibited enhanced growth on PDA supplemented with Cypas® 250 EC at a concentration of 50 ml per 6.25 liters, compared to PDA mixed with Tanidil®-T, which significantly inhibited fungal growth. Spore suspensions obtained from cultures grown on PDA supplemented with Cypas®250 EC demonstrated insect pest control effectiveness comparable to those cultured on PDA alone, with no statistically significant differences observed. This study validates the potential of Cypermethrin, a synthetic pyrethroid, to be effectively combined with the fungus *Metarhizium* spp. for controlling house flies. This approach establishes a foundation for utilizing Cypermethrin in conjunction with other microorganisms to manage various insect species in the future.

ACKNOWLEDGEMENTS

This research project is supported by the Rajamangala University of Technology Isan. Contract No. NKR2562REV023. We are grateful to Ms. Martha Maloi Eromine for editing the language in our manuscript.

AUTHORS' CONTRIBUTIONS

Duanpen Wongsorn: Research project leader, researcher, experimental design, solve problems, write and proof manuscript.

Benya Saenmahayak: researcher, experimental design, solve problems

Nittaya Pitiwittayakul: researcher, experimental design, solve problems

Surachai Rattanasuk: researcher, experimental design, solve problems, write and proof manuscript.

REFERENCES

Abbas M N, Sajeel M and Kausar S. (2013). House fly (*Musca domestica*), a challenging pest; biology, management and control strategies. *Elixir Entomology* 64: 19333–19338. <https://www.researchgate.net/publication/260285055>

- Abbasi E, Yazdani Z, Daliri S and Moemenbellah-Fard M D. (2023). Organochlorine knockdown-resistance (kdr) association in housefly (*Musca domestica*): A systematic review and meta-analysis. *Parasite Epidemiol Control* 22(2023): e00310. <https://doi.org/10.1016/j.parepi.2023.e00310>
- Abbott W S. (1925). A method of computing the effectiveness of insecticides. *Journal of Economic Entomology* 18: 265–267. <https://doi.org/10.1093/jee/18.2.265a>
- Abd El-Ghany T M and Masmali I A. (2016). Fungal biodegradation of organophosphorus insecticides and their impact on soil microbial population. *International Journal of Plant Pathology and Microbiology* 7: 5. <https://doi.org/10.4172/2157-7471.1000349>
- Abidin A F, Ekowati N and Ratnaningtyas N I. (2017). Insecticide compatibility to the entomopathogenic fungi *Beauveria bassiana* and *Metarhizium anisopliae*. *Scripta Biological* 4(4): 273–279. <https://doi.org/10.20884/1.sb.2017.4.4.695>
- Acevedo G R, Zapater M and Toloza A C. (2009). Insecticide resistance of house fly, *Musca domestica* (L.) from Argentina. *Parasitology Research* 105: 489–493. <https://doi.org/10.1007/s00436-009-1425-x>
- Akbar S, Freed, S, Hameed A, Gul H T, Akmal M, Malik M N, Naeem M and Khan M B. (2012). Compatibility of *Metarhizium anisopliae* with different insecticides and fungicides. *African Journal of Microbiology Research* 6(17): 3956–3962. <https://doi.org/10.5897/AJMR12.417>
- Ali Md H, Lailatul F, Md Eunus A S, Mohajira B, Mehenaz S, Md Sabir H, Farha M J, Md. Saiful I F, Supriya A and Md. Selim R. (2024). Exploring the impact of culture media on life history characteristics of the house fly, *Musca domestica* L. (Diptera: Muscidae). *Scientific African* 24: e02133. <https://doi.org/10.1016/j.sciaf.2024.e02133>
- Amutha M, Banu J G, Surulivelu T and Gopalakrishnan N. (2010). Effect of commonly used insecticides on the growth of white muscardine fungus, *Beauveria bassiana* under laboratory conditions. *Journal of Biopesticides* 3(1): 143–146. <https://www.researchgate.net/publication/259357209>
- Antonio B F, Almeida J E M and Clovis L. (2001). Effect of Thiamethoxam on entomopathogenic microorganisms. *Neotropical Entomology* 30(3): 437–447. <https://doi.org/10.1590/S1519-566X2001000300017>
- Apoorva S and Ramaswamy K. (2013). Compatibility of entomopathogenic fungi *Metarhizium anisopliae* (Ascomycota: Hypocreales) with few Pyrethroid and Organophosphate pesticides. *Journal of Pure and Applied Microbiology* 7(1): 721–726. <https://doi.org/10.2478/v10045-010-0004-6>
- Balachander M, Remadevi O K, Sasidharan T O and Bai N S. (2009). Infectivity of *Metarhizium anisopliae* (Deuteromycotina: Hyphomycetes) isolates to the arboreal

- termite *Odontotermes* sp. (Isoptera: Termitidae). *International Journal of Tropical Insect Science* 29: 202–207. <https://doi.org/10.1017/S1742758409990294>
- Barata C, Solayan A and Porte C. (2004). Role of B-esterases in assessing toxicity of organophosphorus (chlorpyrifos, malathion) and carbamate (carbofuran) pesticides to *Daphnia magna*. *Aquatic Toxicology* 66: 125–139. <https://doi.org/10.1016/j.aquatox.2003.07.004>
- Bihal R. Al-Khayri J M, Banu A N, Kudesia N, Ahmed F K, Sarkar R, Arora A and Abd-Elsalam K A. (2023). Entomopathogenic fungi: An eco-friendly synthesis of sustainable nanoparticles and their nanopesticide properties. *Microorganisms* 11: 1617. <https://doi.org/10.3390/microorganisms11061617>
- Bharathi N S, Pannerselvam M, Krishnaswamy S, Shanmugam A and Thattante P R. (2022). Pathogenic potential of *Metarhizium anisopliae* and *Lecanicillium longisporum* on tea mosquito bug, *Helopeltis theivora* Waterhouse (Hemiptera: Miridae). *The Journal of Basic and Applied Zoology* 83: 33. <https://doi.org/10.1186/s41936-022-00297-4>
- Bugti G A, Bin W, Memon S A, Khaliq G and Jaffar M A. (2020). Entomopathogenic fungi: factors involved in successful microbial control of insect pests. *European Journal of Entomology* 17: 74–83. <https://doi.org/10.3923/je.2020.74.83>
- Elliott M. (1980). Established pyrethroid insecticides. *Journal of Pesticide Science* 11: 119–128. <https://doi.org/10.1002/ps.2780110204>
- Farooq M and Freed S. (2016). Combined effects of *Beauveria bassiana* (Hypocreales: Clavicipitaceae) and insecticide mixtures on biological parameters of *Musca domestica* (Diptera: Muscidae). *Pakistan Journal of Zoology* 48(5): 1465–1476. [http://www.zsp.com.pk/pdf48/1465-1476%20\(33\)%20QPJZ-0419-2015-F%201-7-16%20Revised1.pdf](http://www.zsp.com.pk/pdf48/1465-1476%20(33)%20QPJZ-0419-2015-F%201-7-16%20Revised1.pdf)
- Farooq M and Freed, S. (2016). Infectivity of housefly, *Musca domestica* (Diptera: Muscidae) to different entomopathogenic fungi. *Brazilian Journal of Microbiology* 47(4): 807–816. <https://doi.org/10.1016/j.bjm.2016.06.002>
- Geden C J, Nayduch D, Scott J G, Burgess E R, Gerry A C, Kaufman P E, Thomson J, Pickens V and Machtinger E T. (2021). House fly (Diptera: Muscidae): biology, pest status, current management prospects, and research needs. *Journal of Integrated Pest Management* 2(1): 1–38. <https://doi.org/10.1093/jipm/pmaa021>
- Hiromori H and Nishigaki J. (2001). Factor analysis of synergistic effect between the entomopathogenic fungus *Metarhizium anisopliae* and synthetic insecticides. *Applied Entomology and Zoology* 36: 231–236. <https://doi.org/10.1303/aez.2001.231>
- Iqbal W, Malik M F, Sarwar M K, Azam I, Iram N and Rashda A. (2014). Role of housefly (*Musca domestica*, Diptera; Muscidae) as a disease vector: A review. *Journal of*

- Entomology and Zoology Studies* 2(2): 159–163.
<https://www.researchgate.net/publication/275100264>
- Jantanapim J, Sirimungkararat S and Saksirirat S. (2015). Effect of pesticides on growth and efficacy of *Beauveria bassiana* (Balsamo) Vuillemin for controlling of cassava pink mealybug. *KKU Science Journal* 43(3): 469–479. <https://ph01.tci-thaijo.org/index.php/KKUSciJ/article/view/249396> (In Thai)
- Kumar, A, Mukherjee P, Guha A, Adyantaya S D, Mandale A B, Kumar R and Sastry M. (2000). Amphoterization of colloidal gold particles by capping with valine molecules and their phase transfer from water to toluene by electrostatic coordination with fatty amine molecules. *Langmuir* 16(25): 9775–9783. <https://doi.org/10.1021/la000886k>
- Miana U, Galadima I, Gambo F and Zakaria D. (2018). A review on the use of entomopathogenic fungi in the management of insect pests of field crops. *Journal of Entomology and Zoology Studies* 6(1): 27–32.
<https://www.researchgate.net/publication/323445876>
- Mishra S, Kumar P and Malik A. (2011). Adulticidal and larvicidal activity of *Beauveria bassiana* and *Metarhizium anisopliae* against house fly, *Musca domestica* (Diptera: Muscidae), in laboratory and simulated field bioassays. *Parasitology Research* 108: 1483–1492. <https://doi.org/10.1007/s00436-010-2203-5>
- Mishra, S, Kumar P, Malik A and Satya S. (2011). Adulticidal and larvicidal activity of *Beauveria bassiana* and *Metarhizium anisopliae* against housefly, *Musca domestica* (Diptera: Muscidae), in laboratory and simulated field bioassays. *Parasitology Research* 108: 1483–1492. <https://doi.org/10.1007/s00436-010-2203-5>
- Mohammed A A. (2018). Evaluation the efficacy of *Metarhizium anisopliae* and some plant extracts in controlling the housefly, *Musca domestica* in laboratory conditions. *Kufa Journal for Agricultural Science* 10: 1–17. <https://doi.org/10.21276/kjas>
- Muthabathula P and Biruduganti S. (2022). Analysis of biodegradation of the synthetic pyrethroid cypermethrin by *Beauveria bassiana*. *Current Microbiology* 79: 46. <https://doi.org/10.1007/s00284-021-02744-x>
- Nawaz A, Razzaq F, Razzaq A, Gogi M D, Fernández-Grandon G M, Tayib M, Ayub M A, Sufyan M, Shahid M R, Qayyum M, Naveed M, Ijaz A and Arif M J. (2022). Compatibility and synergistic interactions of fungi, *Metarhizium anisopliae*, and insecticide combinations against the cotton aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae). *Scientific Reports* 12: 4843. <https://doi.org/10.1038/s41598-022-08841-6>
- Olagunju E A. (2022). Housefly: Common zoonotic diseases transmitted and control. *Journal of Zoonotic Diseases* 6(1): 1–10. <https://doi.org/10.22034/jzd.2022.14378>

- Oliveira C N de, Neves P M O J and Kawazoe L S. (2003). Compatibility between the entomopathogenic fungus *Beauveria bassiana* and insecticides used in coffee plantations. *Scientia Agricola* 60(4): 663–667. <https://doi.org/10.1590/S0103-90162003000400009>
- Oliveira D G P, Cardoso R R, Mamprim A P and Angeli L F. (2016). Laboratory and field evaluation of a cypermethrin based insecticide for the control of *Alphitobius Diaperinus* Panzer (Coleoptera: Tenebrionidae) and its in-vitro effects on *Beauveria Bassiana* Bals. Vuill. (Hypocreales: Cordycipitaceae). *Revista Brasileira de Cienciaicola* 18(3): 371–379. <https://doi.org/10.1590/1806-9061-2015-0115>
- Ong S Q, Ahmad H, Ab Majid A H and Jaal Z. (2019). Conservation of agricultural soil using entomopathogenic fungi: An agent with insecticides degradation potential. *IOP Conference Series: Earth and Environmental Science* 380: 012014. <https://doi.org/10.1088/1755-1315/380/1/012014>
- Ong S-Q, Ahmad H, Ab Majid A H and Jaal Z. (2017). Interaction between *Metarhizium anisopliae* (Met.) and the insecticides used for controlling house fly (Diptera: Muscidae) in poultry farm of Malaysia. *Journal of Medical Entomology* 54(6): 1626–1632. <https://doi.org/10.1093/jme/tjx128>
- Pelizza S A, Schalamuk S, Simón M R, Stenglein S A, Pacheco-Marinof S G and Scorsetti C. (2018). Compatibility of chemical insecticides and entomopathogenic fungi for control of soybean defoliating pest, *Rachiplusia nu*. *Revista Argentina de Microbiología* 50(2): 189–201. <https://doi.org/10.1016/j.ram.2017.06.002>
- Ponganan K, Sirimungkararat S and Saksirirat W. (2015). Effect of stickers, UV protectant and common pesticides on survival and efficiency of *Beauveria bassiana* for control brown planthopper. *Khon Kaen Agriculture Journal* 43(Suppl. 1): 323–331. <https://ag2.kku.ac.th/kaj/PDF.cfm?filename=Vol43suppl.1.pdf&id=1787&keeptrack=12>
- Quesada-Moraga E, González-Mas N, Yousef-Yousef M, Garrido-Jurado I and Fernández-Bravo M. (2024). Key role of environmental competence in successful use of entomopathogenic fungi in microbial pest control. *Journal of Pest Science*. 97: 1–15. <https://doi.org/10.1007/s10340-023-01622-8>
- Reynolds J A. (2017) Epigenetic influences on diapause. *Advances in Insect Physiology* 53: 115–144. <https://doi.org/10.1016/bs.aiip.2017.03.003>
- Sain S M, Monga D, Kumar R, Nagrale D T, Hiremani N S and Kranthi S. (2019). Compatibility of entomopathogenic fungi with insecticides and their efficacy for IPM of *Bemisia tabaci* in cotton. *Journal of Pest Science* 44(2): 97–105. <https://doi.org/10.1584/jpestics.D18-067>

- Saranraj P and Jayaprakash A. (2017). Agrobeneficial entomopathogenic fungi – *Beauveria bassiana*: A review. *IAJMR* 3(2): 1051–1087. <https://doi.org/10.22192/iajmr.2017.3.2.4>
- Schumacher V and Poehling H-M. (2012). In vitro effect of pesticides on the germination, vegetative growth, and conidial production of two strains of *Metarhizium anisopliae*. *Fungal Biology* 116: 121–132. <https://doi.org/10.1016/j.funbio.2011.10.007>
- Sharififard M, Mossadegh M S, Vazirianzadeh B and Zarei-Mahmoudabadi A. (2011). Interactions between entomopathogenic fungus, *Metarhizium anisopliae* and sublethal doses of Spinosad for control of house fly, *Musca domestica*. *Journal of Arthropod-Borne Diseases* 5(1): 28–36. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3385566/pdf/ijad-5-28.pdf>
- Sharififard M, Mossadegh M S, Vazirianzadhe B and Mahmoudabadi A Z. (2011). Laboratory pathogenicity of entomopathogenic fungi, *Beauveria bassiana* (Bals.) Vuill. and *Metarhizium anisoplae* (Metch.) Sorok. to larvae and adult of house fly, *Musca domestica* L. (Diptera: Muscidae). *Asian Journal of Biological Sciences* 4: 128–137. <https://scialert.net/abstract/?doi=ajbs.2011.128.137>
- Tamai M A, Alves S B, Lopes R B, Faion M and Padulla L F L. (2002). Toxicidade de produtos fitossanitários para *Beauveria bassiana* (Bals.) Vuill. *Arquivos do Instituto Biologico* 69: 89–96. http://www.biologico.sp.gov.br/uploads/docs/arq/V69_3/Tamai1.pdf <https://doi.org/10.1590/1808-1657v69n3p0892002>
- Wang J-N, Hou J, Wu Y-Y, Guo S, Liu Q-M, Li T-Q and Gong Z-Y. (2019). Resistance of house fly, *Musca domestica* L. (Diptera: Muscidae), to five insecticides in Zhejiang Province, China: the situation in 2017. *Canadian Journal of Infectious Diseases* 4851914. <https://doi.org/10.1155/2019/4851914>