



Caffeine Prolongs Survivability and Significantly Modulates Cytokine Levels of TNF- α , IFN- γ , IL-18 and IL-10 in Plasmodium berghei ANKA-infected ICR Mice

Authors:

Ahmad Zakwan Noriman, Mohd Ridzuan Mohd Abd Razak, Wan Nurhayati Wan Hanafi, Rusliza Basir, Hani Kartini Agustar and Wan Rozianoor Mohd Hassan*

***Correspondence:** rozianoor@uitm.edu.my

Submitted: 21 April 2025; **Accepted:** 15 January 2026; **Published:** 31 July 2026

To cite this article: Ahmad Zakwan Noriman, Mohd Ridzuan Mohd Abd Razak, Wan Nurhayati Wan Hanafi, Rusliza Basir, Hani Kartini Agustar and Wan Rozianoor Mohd Hassan (2026). Caffeine prolongs survivability and significantly modulates cytokine levels of TNF- α , IFN- γ , IL-18 and IL-10 in *Plasmodium berghei* ANKA-infected ICR mice. *Tropical Life Sciences Research* 37(2): 25–52. <https://doi.org/10.21315/tlsr2026.37.2.2>

To link to this article: <https://doi.org/10.21315/tlsr2026.37.2.2>

HIGHLIGHTS

- The lowest concentration of caffeine (5 mg/kg b.w.) showed the most effective chemosuppression at $64.18 \pm 9.17\%$ as compared to other concentrations (10 mg/kg b.w and 20 mg/kg b.w.) in *Plasmodium berghei* ANKA-infected ICR mice.
- Caffeine at 5 mg/kg b.w. prolongs the survivability (17 days) of *P. berghei* ANKA-infected ICR mice.
- Caffeine significantly modulates levels of pro- and anti-inflammatory cytokines (TNF- α , IFN- γ , IL-18 and IL-10) in cerebral malaria murine model.

Caffeine Prolongs Survivability and Significantly Modulates Cytokine Levels of TNF- α , IFN- γ , IL-18 and IL-10 in *Plasmodium berghei* ANKA-infected ICR Mice

¹Ahmad Zakwan Noriman, ²Mohd Ridzuan Mohd Abdul Razak, ^{1,3}Wan Nurhayati Wan Hanafi, ⁴Rusliza Basir, ⁵Hani Kartini Agustar and ^{1,3}Wan Rozianoor Mohd Hassan*

¹School of Biology, Faculty of Applied Sciences, Universiti Teknologi MARA, 40450 Shah Alam, Selangor, Malaysia

²Bioassay Unit, Herbal Medicine Research Center, Institute for Medical Research, National Institute of Health, 40170 Shah Alam, Selangor, Malaysia

³Human Genetic and Biochemistry, Research Nexus of UiTM, Universiti Teknologi MARA, 40450 Shah Alam, Selangor, Malaysia

⁴Pharmacology Unit, Department of Human Anatomy, Faculty of Medicine and Health Science, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

⁵Department of Earth Science and Environment, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

Submitted: 21 April 2025; **Accepted:** 15 January 2026; **Published:** 31 July 2026

To cite this article: Ahmad Zakwan Noriman, Mohd Ridzuan Mohd Abd Razak, Wan Nurhayati Wan Hanafi, Rusliza Basir, Hani Kartini Agustar and Wan Rozianoor Mohd Hassan (2026). Caffeine prolongs survivability and significantly modulates cytokine levels of TNF- α , IFN- γ , IL-18 and IL-10 in *Plasmodium berghei* ANKA-infected ICR mice. *Tropical Life Sciences Research* 37(2): 25–52. <https://doi.org/10.21315/tlsr2026.37.2.2>

To link to this article: <https://doi.org/10.21315/tlsr2026.37.2.2>

Abstract: Caffeine, a bioactive compound in *Theobroma cacao*, exhibits various pharmacological activities, including anti-inflammatory properties. Malaria is an inflammatory-related disease caused by *Plasmodium* spp. Severe infection of this disease can cause cerebral malaria, which is characterised by neuroinflammation. An *in vitro* study of caffeine suggested that it possesses anti-malarial properties. However, caffeine's potential in both anti-malarial and cytokine-modulating properties through *in vivo* have yet to be explored. This study investigated the anti-malarial and cytokine-modulating effects of caffeine in *P. berghei* ANKA-infected ICR mice. Anti-malarial activity of caffeine at 5 mg/kg, 10 mg/kg and 20 mg/kg body weight was given via intraperitoneal (i.p.) injection, and the chemosuppressive percentage (percentage of parasitaemia inhibition) was calculated based on Peter and Robinsons's study in 1992. In addition, the cytokine-modulating effect was determined using the ELISA method. Chemosuppressive test at 5 mg/kg, 10 mg/kg and 20 mg/kg b.w. of caffeine in the *P. berghei* ANKA-infected ICR mice exhibited 64.18 ± 9.17 , 39.93 ± 5.63 and $21.86 \pm 5.57\%$, respectively. Notably, the lowest concentration, 5 mg/kg b.w. of caffeine was able to prolong significantly ($p < 0.05$) the survivability of malaria-infected ICR mice for 17 days as compared to infected mice (12 days). This effective dose of caffeine revealed a significant reduction in cytokine levels of TNF- α , IFN- γ , IL-18 and IL-10 by 1.08-, 1.12-, 1.12- and 1.15-folds, respectively, as compared to malaria-infected mice. Collectively, 5 mg/kg b.w. of caffeine exhibited good chemosuppressive, modulated levels of inflammatory cytokines and prolonged survivability of *P. berghei* ANKA-infected ICR mice. Hence, caffeine can be a plausible candidate for the development of anti-malarial therapeutic drugs.

Keywords: Caffeine, Cytokine-Modulation, ICR Mice, Malaria, *P. berghei* ANKA

*Corresponding author: rozianoor@uitm.edu.my

Abstrak: Kafein, sebatian bioaktif dalam *Theobroma cacao*, mempamerkan pelbagai aktiviti farmakologi, termasuk sifat anti-radang. Malaria merupakan penyakit berkaitan keradangan yang disebabkan oleh *Plasmodium* spp. Jangkitan ketara penyakit ini boleh menyebabkan malaria serebrum, yang dicirikan oleh keradangan saraf. Kajian in vitro kafein mencadangkan bahawa ia mempunyai kandungan anti-malaria. Walau bagaimanapun, potensi kafein yang mempunyai aktiviti anti-malaria dan modulasi sitokin dalam in vivo masih belum diterokai. Kajian ini bertujuan untuk mengenal pasti kesan anti-malaria dan modulasi sitokin kafein dalam mencit ICR yang dijangkiti *P. berghei* ANKA. Aktiviti anti-malaria kafein pada 5 mg/kg, 10 mg/kg dan 20 mg/kg berat badan (b.b.) diberikan melalui suntikan intraperitoneum (i.p.) dan peratusan kemosupresif (peratusan perencatan parasitemia) dikira berdasarkan kajian Peter dan Robinsons pada tahun 1992. Tambahan pula, kesan modulasi sitokin ditentukan menggunakan kaedah ELISA. Ujian kemosupresif pada 5 mg/kg, 10 mg/kg dan 20 mg/kg b.b. kafein dalam mencit ICR yang dijangkiti *P. berghei* ANKA menunjukkan 64.18 ± 9.17 , 39.93 ± 5.63 , dan $21.86 \pm 5.57\%$ masing-masing. Khususnya, kafein dengan kepekatan terendah, 5 mg/kg b.b. dapat memanjangkan dengan ketara ($p < 0.05$) kemandirian mencit ICR yang dijangkiti malaria selama 17 hari berbanding dengan mencit yang dijangkiti (12 hari). Rawatan dengan dos kafein yang berkesan ini mendedahkan pengurangan ketara dalam aras sitokin TNF- α , IFN- γ , IL-18 dan IL-10 berbanding mencit yang dijangkiti malaria. Secara kolektif, kafein pada 5 mg/kg b.b. mempamerkan penindasan kemosupresif yang baik, memodulasi aras keradangan sitokin dan memanjangkan kemandirian mencit ICR yang dijangkiti *P. berghei* ANKA. Oleh itu, kafein boleh menjadi calon pembangunan ubat terapeutik anti-malaria.

Kata kunci: Kafein, Modulasi Sitokin, Mencit ICR, Malaria, *P. berghei* ANKA

INTRODUCTION

Malaria is an infectious disease globally, as 95% of the diseases accounted for are from major regions of African countries (World Health Organization 2024). In 2023, 263 million infections and 597,000 deaths worldwide were reported due to this inflammatory-related disease (World Health Organization 2024). *P. knowlesi*, *P. ovale*, *P. falciparum*, *P. vivax* and *P. malariae* are parasite species that can infect humans (Lee et al. 2022). A severe complication of malaria infection is cerebral malaria, which is characterised by the sequestration of parasitised red blood cells (pRBC) within the microcirculation and elevated pro-inflammatory cytokines such as IFN- γ , TNF- α and IL-18 (Ali et al. 2022; Basir et al. 2012).

Chloroquine (CQ), a common malaria treatment, is synthesised from quinoline that was found in Cinchona bark in the early 19th century (Zhou et al. 2020). The emergence and spread of drug-resistant *Plasmodium* parasites, such as CQ and artemisinin, can also contribute to the increase in malaria cases (Parija & Antony 2016). Rising cases of anti-malarial drug resistance prompted the need to develop and improve drugs for treatment, especially from medicinal plants (Rasmussen et al. 2022; Titanji et al. 2008; Vaou et al. 2021). Many anti-malarial drug development studies have been conducted from plant resources (Habibi et al. 2022). Latest research on malaria includes exploration of natural plants such

as the methanol extract of *Picralima nitida* in Swiss albino mice infected with *P. berghei* and the ethanol extract of *Blighia sapida* K.D. Koenig in Swiss albino mice infected with CQ-resistant *P. berghei* ANKA MRA 311 (Ogbeiden et al. 2023; Akinnyemi et al. 2025).

1,3,7-trimethylxanthine, or caffeine, is a major bioactive compound that is consumed daily by humans in the form of coffee, tea, or chocolate (Dorvigny et al. 2022; Temple et al. 2017). Caffeine has been known as a stimulant in the central nervous system (Chen et al. 2010). Besides that, caffeine has been identified as having protective effects against neurodegenerative diseases by preventing and stabilising the blood-brain barrier to ensure that the barrier remains intact (Chen et al. 2010). The alkaloid has also shown anti-inflammatory and neuroprotective abilities (Barcelos et al. 2020; Kolahehdouzan & Hamadeh 2017). In addition, caffeine is an A2AR antagonist which competes with adenosine to bind to the receptor. Caffeine binding to the A2AR receptor inhibited the binding of adenosine, which caused the A2AR not to be activated. This further blocked NF- κ B, which reduces inflammatory cytokine production and prevents neuroinflammation (Sonsalla et al. 2012).

Caffeine at 20 mg/mL pyrazolines has been identified to have potential for anti-malarial properties based on an *in vitro* study with a high inhibition percentage ($85.2 \pm 5.4\%$) against *P. falciparum* (Insuasty et al. 2015). However, caffeine's potential as an anti-malarial property and cytokine-modulating effect through *in vivo* model has yet to be explored. A previous study by Basir et al. (2012) was able to mimic the *P. falciparum* infection in humans using *Plasmodium berghei* ANKA in ICR mice (Basir et al. 2012). *P. berghei* ANKA is a widely used rodent model to study cerebral malaria, particularly in the context of evaluating anti-malarial properties and understanding immune responses (Craig et al. 2012). Thus, this study aims to investigate the anti-malarial and cytokine-modulating effects of caffeine in *P. berghei* ANKA-infected ICR mice.

MATERIALS AND METHODS

Parasite

P. berghei ANKA strain (CQ-sensitive) was procured from The Bioassay Unit, National Institute of Health (NIH), Shah Alam, Malaysia. A combination of cryogenic storage and passage in ICR mice was done to conserve the parasite at Universiti Teknologi MARA (UiTM) Shah Alam, Malaysia.

Treatment

Caffeine (1,3,7-trimethylxanthine; CAS No. 58-08-2) with a purity of 99.99% was purchased from Sigma-Aldrich and diluted in distilled water to prepare treatment concentrations of 5 mg/kg, 10 mg/kg and 20 mg/kg b.w. The treatments were administered via i.p. injection. For the positive control, CQ (CAS No. 50-63-5), also obtained from Sigma-Aldrich at a purity of 99.99%, was diluted in distilled water and administered at 10 mg/kg b.w.

Experimental Mice

Male ICR mice weighing between 20 g and 25 g were procured from Laboratory Animal Facility and Management (LAFAM), UiTM Puncak Alam, and were accommodated at Animal Lab House, UiTM Shah Alam. International guidelines for the use and maintenance of experimental animals were used for the maintenance and care of the mice. The rules and regulations set by the UiTM Committee on Animal Research and Ethics (UiTM CARE) of UiTM Shah Alam were followed for all procedures conducted on the animals (Ethical approval number: UiTM CARE: 399/2023 [3 February 2023]).

Sample Size Calculation

In the experiment, a 10% dropout rate was considered to account for potential intraperitoneal (IP) injection failures, as reported by Das and North (2007). The likelihood of unsuccessful parasite infection was not included in the calculation, as the probability of mice failing to become infected is less than 1% (Leitner *et al.* 2010). According to Basir *et al.* (2012), mortality in untreated infected mice increases only after the treatment period, with rates of 20% on day 5, 70% on day 6 and 100% by day 7. During the treatment procedure, the survival rate is typically 100%, as deaths occur after treatment is completed. Nonetheless, to accommodate the small possibility that a mouse might die during the 4-day treatment phase, a precautionary 5% mortality rate was included in the calculation.

Survivability test (Ali et al. 2017; 2022; Hassan et al. 2019)

A total of six groups were used in the study, each consisting of seven mice ($n = 7$), including a normal control, an infected control, a CQ-treated group (10 mg/kg) and three caffeine-treated groups (5 mg/kg, 10 mg/kg and 20 mg/kg), requiring 42 male ICR mice. An additional four mice were needed for parasite culture. Allowances were also made for potential

losses, including a 5% mortality rate among infected mice (2 mice) and a 10% IP injection failure rate (5 mice). In total, 53 male ICR mice aged 6 weeks–8 weeks were required for the study.

Cytokine Modulation activity (Ali et al. 2017; 2022; Hassan et al. 2019)

In this study, a minimum of 40 male mice was required, consisting of four groups with 10 mice each: a normal control group, an infected group, an infected group treated with CQ and an infected group treated with the effective dose of caffeine. An additional three mice were used for parasite culture before group inoculation. To account for potential losses, a 5% mortality rate was considered for the 30 infected mice, adding approximately two mice, and a 10% IP injection failure rate was applied to the 40 mice receiving injections, adding another four mice. Therefore, a total of 49 male mice were required for the study.

Anti-Malarial Assay

Inoculum preparation

The serial passage of *P. berghei* was initiated based on a previous study (Huang et al. 2015). Stored blood parasites in Alservers' buffer solution were IP injected into normal mice at 0.2 mL and allowed for three to five days of inoculation. After being anaesthetised with carbon dioxide, a cardiac puncture was done to draw blood from infected mice. A volume of 0.2 mL containing 2×10^7 pRBC was injected into a normal mouse as a subsequent passage. Blood was diluted with saline 0.85% to give 2×10^7 pRBC in 0.2 mL of injection volume, which accounted for approximately 50% of the percentage of parasitaemia (Basir et al. 2012).

Anti-Malarial Activity (4-Day Suppressive Test)

Male ICR mice ($N = 42$) were distributed into six groups of seven mice ($n = 7$), following simple randomisation (Verhave et al. 2024) and each was given a different treatment (Ali et al. 2017; 2022; Hassan et al. 2019). The groups of mice were the normal (saline 0.85%), *P. berghei* ANKA-infected (saline 0.85%), CQ-treated infected (10 mg/kg b.w.) and caffeine-treated infected (5 mg/kg, 10 mg/kg and 20 mg/kg b.w.). For the malarial infection, the mice were injected with 0.2 mL of infected blood (2×10^7 pRBC) for four consecutive days (4-day suppressive test) before treatments.

Throughout the study, the basic parameters such as physical changes, body weight and body temperature were measured and recorded daily. The body weight of the mice was measured using a weighing balance

(Camry, Hong Kong), while the axillary temperature was measured using a thermometer (medACCU, China). All mice were visually observed for signs of illness or behavioural changes such as lethargy, decreased locomotor activity, piloerection and diarrhoea. Any indications of illness were categorised as absent (-), mild (+), moderate (++) or severe (+++) using an arbitrary scale. Survivability of the mice was recorded for 30 days. The thin blood smear was prepared every two days to observe the parasitaemia count.

Parasitaemia Measurement

The staining procedure was conducted based on Huang *et al.* (2015). A thin blood film was developed from the collected blood drop and stained with Giemsa's stain. The blinding procedure was implemented to minimise observer bias. The parasitaemia percentage was calculated in random order of slides from five random sites up to 1,000 cells in total using a light microscope (Leica, Germany). This procedure was conducted to prevent systematic measurement bias that adheres to the principles outlined by the ARRIVE guidelines (Percie du Sert *et al.* 2020; Verhave *et al.* 2024).

The percentage of parasitaemia and chemosuppression was calculated using Peters and Robinson (1992) with slight modification.

Percentage of Parasitaemia = (Number of parasitised RBC/Total number of RBC count) × 100

Percentage of Chemosuppression = ([Parasitemia in negative control – Parasitemia in the study group]/Parasitaemia in negative control) × 100

The animal's mean survival time was calculated using the following formula (Mekonnen 2015):

Mean survival time (MST) = (Total number of mice survived/Total number of mice)

Cytokine Modulation Activity

The suppressive test was repeated with four groups of 10 mice: normal, infected mice, CQ-treated infected (10 mg/kg b.w.) and caffeine-treated infected (5 mg/kg b.w.). Upon completion of treatment (Day 5), all the mice were euthanised, and blood was drawn via cardiac puncture.

The pro-and anti-inflammatory cytokines were measured in the mice's serum by means of the enzyme-linked immunoassay (ELISA) method using the readily accessible kits. The pro-inflammatory cytokines include TNF- α (ELABSCIENCE, USA; Catalogue number: E-EL-M3063; Detection limit: 4.69 pg/mL), IFN- γ (ELABSCIENCE, USA; Catalogue number: E-EL-M0048; Detection limit: 9.38 pg/mL) and IL-18 (MLB International, USA; Catalogue number: 7625; Detection limit 25.0 pg/mL). The anti-inflammatory cytokines include IL-4 (ELABSCIENCE, USA; Catalogue number IL-4: E-EL-M0043; Detection limit: 18.75 pg/mL) and IL-10 (ELABSCIENCE, USA; Catalogue number IL-10: E-EL-M0046; Detection limit: 9.38 pg/mL).

Statistical Analysis

The results were analysed using GraphPad Prism version 10.1.1 and the data presented as mean value $\pm$ SEM. One-way analysis of variance (ANOVA) followed by Tukey's test as a single post-hoc test was used to compare the experiment group means with $p < 0.05$, $p < 0.01$ and $p < 0.001$ considered statistically significant. Student's t-test and log-rank test were used to evaluate the data for Kaplan-Meier survival analysis.

RESULTS

Anti-Malarial Activity (4-Day Suppressive Test)

Mice administered with 5 mg/kg b.w. of caffeine for four consecutive days following *P. berghei* ANKA infection resulted in significantly ($p < 0.05$) lowest parasite percentage of $8.38 \pm 0.35\%$ as compared to infected mice (Table 1; Fig. 1). Chemosuppressive effects of caffeine (5 mg/kg b.w.) demonstrated a significant ($p < 0.05$) increased percentage at $65.47 \pm 1.42\%$ as compared to infected mice (Table 2; Fig. 2). The median survival rate improved significantly in mice treated with 5 mg/kg b.w. of caffeine (17 days) compared to infected mice (12 days) (Table 3; Fig. 3). The CQ-treated infected group exhibited the lowest parasite growth ($1.22 \pm 0.25\%$), the highest chemosuppression (94.99 ± 1.02), and the highest mean survival time (30 days) compared to other treatment groups. This is consistent with its function as an established anti-malarial reference drug (Zhou et al. 2020).

TABLE 1. Parasitised red blood cells percentage (pRBC%) of mice infected with *P. berghei* ANKA with or without caffeine treatment on Day-4.

Group	Normal (n = 7)	Infected mice (negative control) (n = 7)	CQ-treated (10 mg /kg b.w.) (positive control) (n = 7)	Caffeine treated		
				(5 mg/kg b.w.) (n = 7)	(10 mg/kg b.w.) (n = 7)	(20 mg/ kg b.w.) (n = 7)
Mean PRBC percentage (%)	0	24.26 ± 5.11 ^b ^{b-p < 0.001}	1.22 ± 0.25 ^{a-p} < 0.001	8.38 ± 0.35 ^{a-p < 0.001,} ^{b-p = 0.013}	14.90 ± 0.33 ^{b-p = 0.003}	18.96 ± 0.49 ^{b-p} = 0.01

Notes: Mean PRBC was calculated on day-4 post infection. Data represent mean ± S.E.M for mean pRBC percentage (n = 7). A = Significantly different comparison with infected control; b = Significantly different comparison with CQ-treated; b.w. = body weight; CQ = chloroquine; PRBC = parasitised red blood cells.


**FIGURE 1:** Parasitised red blood cell (pRBC) percentage of mice infected with *P. berghei* ANKA with or without caffeine treatment throughout 30 days of monitoring. Data represent the pRBC percentage of normal (non-infected mice) (n = 7), negative control (nontreated *P. berghei* ANKA-infected mice) (n = 7), CQ-treated mice (n = 7), caffeine-treated (5 mg/kg b.w.) (n = 7), caffeine-treated (10 mg/kg b.w.) (n = 7) and caffeine-treated (20 mg/kg b.w.) (n = 7). A significant difference between groups was determined at **p* < 0.05, ***p* < 0.01, ****p* < 0.001.

TABLE 2. Chemosuppression of mice infected with *P. berghei* anka with or without caffeine treatment on day-4.

Group	Normal (n = 7)	Infected mice (negative control) (n = 7)	CQ-treated (10 mg /kg b.w.) (positive control) (n = 7)	Caffeine treated		
				(5 mg/kg b.w.) (n = 7)	(10 mg/kg b.w.) (n = 7)	(20 mg/ kg b.w.) (n = 7)
Chemosuppression	100	0	94.99 ± 1.02 ^{a-} P<0.001	65.47 ± 1.42 ^{a-} P<0.001, b-P<0.001	38.57 ± 1.38 ^{a-} P<0.001, b-P<0.001	21.85 ± 2.01 ^{a-} P<0.001, b-P<0.001

Notes: Chemo-suppression was calculated on day-4 post infection. Data represent mean ± S.E.M for chemosuppression (n = 7); a = Significantly different from infected control comparison; b = Significantly different from CQ-treatment comparison; b.w. = body weight; CQ = chloroquine.





FIGURE 2: Chemosuppressive of mice infected with *P. berghei* ANKA with or without caffeine treatment throughout 30 days of monitoring. Data represent chemosuppression of normal (non-infected mice) (n = 7), negative control (non-treated *P. berghei* ANKA-infected mice) (n = 7), CQ-treated mice (n = 7), caffeine-treated (5 mg/kg b.w.) (n = 7), caffeine-treated (10 mg/kg b.w.) (n = 7) and caffeine-treated (20 mg/kg b.w.) (n = 7). A significant difference between groups was determined at **p* < 0.05, ***p* < 0.01, ****p* < 0.001.

TABLE 3. Median survival time of mice infected with *P. berghei* ANKA with or without caffeine treatment throughout 30 days of monitoring.

Group	Normal (<i>n</i> = 7)	Infected mice (negative control) (<i>n</i> = 7)	CQ-treated (10 mg /kg b.w.) (positive control) (<i>n</i> = 7)	Caffeine treated		
				(5 mg/kg b.w.) (<i>n</i> = 7)	(10 mg/kg b.w.) (<i>n</i> = 7)	(20 mg/ kg b.w.) (<i>n</i> = 7)
Mean survival time	30 days a- <i>p</i> < 0.001	12 days ^{b-p} < 0.001	30 days ^{a-p} < 0.001	17 days ^{a-p} ; ^{b-p} < 0.001	15 days ^{b-p} < 0.001	11 days ^{b-p} < 0.001

Notes: Median survival time was calculated on day-30 post infection. Data represent mean $\pm$ S.E.M for survival (*n* = 7). a = Significantly different comparison with infected control; b = Significantly different comparison with CQ-treated; b.w. = body weight; CQ = chloroquine.

**FIGURE 3:** Representative Kaplan-Meier survival curve of mice infected with *P. berghei* ANKA with or without caffeine treatment in a 30-day survivability test. Data represent survival of normal (non-infected mice) (*n* = 7), negative control (non-treated *P. berghei* ANKA-infected mice) (*n* = 7), CQ-treated mice (*n* = 7), caffeine-treated (5 mg/kg b.w.) (*n* = 7), caffeine-treated (10 mg/kg b.w.) (*n* = 7) and caffeine-treated (20 mg/kg b.w.) (*n* = 7). A significant difference between groups was determined at **p* < 0.05, ***p* < 0.01 and ****p* < 0.001.

The mice body weight in all groups was not significantly different (*p* > 0.05) when compared to normal mice (Fig. 4; Table 4). The body weight of the mice slightly increased for the normal group and CQ-treated infected group throughout 30 days of the survival test (Fig. 4). Mice from

the caffeine-treated infected groups exhibited decreased body weight. However, they were not significantly reduced ($p > 0.05$) throughout the 30 days of monitoring after being given treatment for four consecutive days (Fig. 4). In addition, the body temperature of the mice in all groups was not significantly ($p > 0.05$) different from normal (Fig. 5; Table 5). Physical illness, including lethargy, decreased locomotor activity and urine colour changes, were observed at 5 mg/kg b.w. of caffeine much later compared to other caffeine-treated infected groups (Appendix).

TABLE 4. Mean body weight of mice infected with *p. berghei* anka with or without caffeine treatment on day-4.

Group	Normal (n = 7)	Infected mice (negative control) (n = 7)	CQ-treated (10 mg /kg b.w.) (positive control) (n = 7)	Caffeine treated		
				(5 mg/kg b.w.) (n = 7)	(10 mg/kg b.w.) (n = 7)	(20 mg/ kg b.w.) (n = 7)
Body weight (grams)	29.83 ± 1.47 n.d.	29.17 ± 1.17 n.d.	28.83 ± 2.86 n.d.	30.17 ± 4.4 n.d.	31.17 ± 3.251 n.d.	29.5 ± 5.86 n.d.

Notes: Body weight was calculated on day-4 post infection. Data represent mean ± S.E.M for body weight (n = 7). a = Significantly different comparison with infected control; b = Significantly different comparison with CQ-treated; n.d. = No significantly different comparison with control groups; b.w. = body weight; CQ = chloroquine.





FIGURE 4: Body weight of mice infected with *P. berghei* ANKA with or without caffeine treatment throughout 30 days of monitoring. Data represent survival of normal (non-infected mice) (n = 7), negative control (nontreated *P. berghei* ANKA-infected mice) (n = 7), CQ-treated mice (n = 7), caffeine-treated (5 mg/kg b.w.) (n = 7), caffeine-treated (10 mg/kg b.w.) (n = 7) and caffeine-treated (20 mg/kg b.w.) (n = 7). A significant difference between groups was determined at * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$.

TABLE 5. Mean body temperature of mice infected with *P. berghei* ANKA with or without caffeine treatment on day-4.

Group	Normal (n = 7)	Infected mice (negative control) (n = 7)	CQ-treated (10 mg /kg b.w.) (positive control) (n = 7)	Caffeine treated		
				(5 mg/kg b.w.) (n = 7)	(10 mg/kg b.w.) (n = 7)	(20 mg/ kg b.w.) (n = 7)
Body temperature (°C)	37.1 ± 0.17 n.d.	37.08 ± 0.18 n.d.	37.12 ± 0.21 n.d.	37.02 ± 0.23 n.d.	36.85 ± 0.3 n.d.	36.72 ± 0.33 n.d.

Notes: Body temperature was calculated on day-4 post infection. Data represent mean ± S.E.M for body temperature (n = 7). a = Significantly different comparison with infected control; b = Significantly different comparison with CQ-treated; n.d. = No significantly different comparison with control groups; b.w. = body weight; CQ = chloroquine.


**FIGURE 5:** Body temperature of mice infected with *P. berghei* ANKA with or without caffeine treatment throughout 30 days of monitoring. Data represent survival of normal (non-infected mice) (n = 7), negative control (nontreated *P. berghei* ANKA-infected mice) (n = 7), CQ-treated mice (n = 7), caffeine-treated (5 mg/kg b.w.) (n = 7), caffeine-treated (10 mg/kg b.w.) (n = 7) and caffeine-treated (20 mg/kg b.w.) (n = 7). A significant difference between groups was determined at * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Cytokine-Modulation Activity


The serum cytokine levels of the mice were examined for modulations of cytokine activities. The present result in Fig. 6, Fig. 7 and Table 6 indicates that there was an increase in both pro-and anti-inflammatory cytokines

on day-4 after *P. berghei* ANKA infection. Parasite infection increased the level of the pro- and anti-inflammatory cytokines of TNF- α , IL-18, IFN- γ , IL-4 and IL-10 at 9.70-, 4.89-, 47.89-, 2.77- and 2.75-folds, respectively, compared to normal mice. CQ exhibited reductions in TNF- α , IL-18, IFN- γ , IL-4 and IL-10 by 9.41-, 4.37-, 27.01-, 1.2- and 2.39-folds, respectively, compared to infected mice (Table 6). The pro- and anti-inflammatory cytokines of TNF- α , IFN- γ , IL-18 and IL-10 were significantly reduced (by 1.08-, 1.12-, 1.12- and 1.15-folds, respectively) in the serum after four days of caffeine treatment at 5 mg/kg b.w. as compared to the infected mice (Fig. 6; Fig. 7; Table 7). IL-4 cytokine level did not exhibit any significant difference when treated with caffeine compared to the infected mice (1.09-fold). Levels of pro-and anti-inflammatory cytokines of TNF- α , IL-18, IFN- γ and IL-10 were significantly reduced with CQ treatment. CQ was able to enhance parasite infection by reducing inflammatory cytokine production since it can eradicate parasites in the initial phase of infection. Like quinine and mefloquine, CQ can directly affect the immune system (Kwiatkowski & Bate 1995; Picot *et al.* 1993). Although CQ at 10 mg/kg b.w. can significantly reduce IL-4 cytokine level (2.3-fold), the level is not as low as in normal mice (Table 6). The same pattern was exhibited by caffeine treatment, which did not significantly reduce IL-4 cytokine level as compared to infected mice.

(a)



(b)



(c)





FIGURE 6: Serum concentration of pro-inflammatory cytokines of (a) TNF- α ; (b) IFN- γ and (c) IL-18 in normal, infected mice, CQ-treated (10 mg/kg b.w.) and caffeine-treated (5 mg/kg b.w.) on day-4 post-infection. Data represent mean $\pm$ S.E.M of normal ($n = 10$), infected mice ($n = 10$), CQ-treated (10 mg/kg b.w.) ($n = 10$), and caffeine-treated (5 mg/kg b.w.) mice ($n = 10$). A significant difference between groups was determined at $*p < 0.05$, $**p < 0.01$ and $***p < 0.001$. CQ = chloroquine; IFN = Interferon; IL = interleukin; TNF = Tumour Necrosis Factor.

(a)



(b)




FIGURE 7: Serum concentration of anti-inflammatory cytokines of (a) IL-4 and (b) IL-10 with normal, infected mice, CQ treated (10 mg/kg b.w.) and caffeine treated (5 mg/kg b.w.) on day-4 post-infection. Data represent mean $\pm$ S.E.M of normal ($n = 10$), infected mice ($n = 10$), CQ-treated (10 mg/kg b.w.) ($n = 10$), and caffeine-treated (5 mg/kg b.w.) mice ($n = 10$). A significant difference between groups was determined at $*p < 0.05$, $**p < 0.01$ and $***p < 0.001$. CQ = chloroquine; IL = interleukin.

TABLE 6. Inflammatory cytokine levels of mice infected with *P. berghei* ANKA treated with or without caffeine treatment.

Cytokines		Groups			
		Normal (n = 10)	Infected mice (negative control) (n = 10)	CQ-treated (10 mg / kg b.w.) (positive control) (n = 10)	Caffeine (5 mg/kg b.w.) (n = 10)
Pro- inflammatory cytokine	Tumour nuclear factor alpha (TNF- α)	74.54 $\pm$ 1.48 ^{a-P} < 0.001	722.8 $\pm$ 15.27 ^{b-P} < 0.001	76.78 $\pm$ 3.08 a-P < 0.001	670.6 $\pm$ 14.82 ^{a-P} = 0.009, b-P < 0.001
	Interferon Gamma (IFN- γ)	7.733 $\pm$ 0.74 ^{a-P} < 0.001	370.3 $\pm$ 10.30 ^{b-P} < 0.001	13.71 $\pm$ 2.04 ^{a-} P < 0.001	331.3 $\pm$ 13.09 ^{a-P} = 0.013, b-P < 0.001
	Interleukin-18 (IL-18)	73.85 $\pm$ 2.03 ^{a-P} < 0.001	361.0 $\pm$ 13.40 ^{b-P} < 0.001	82.59 $\pm$ 2.57 ^{a-P} < 0.001	322.4 $\pm$ 9.63 ^{a-P} = 0.015, b-P < 0.001
Anti- inflammatory cytokine	Interleukin-4 (IL-4)	13.43 $\pm$ 1.55 ^{a-P} < 0.001	37.22 $\pm$ 1.59 ^{b-P} = 0.019	31.00 $\pm$ 0.69 a-P = 0.019	34.16 $\pm$ 1.55 n.d.
	Interleukin-10 (IL-10)	159.7 $\pm$ 12.49 ^{a-P} < 0.001	439.6 $\pm$ 16.10 ^{b-P} < 0.001	174.7 $\pm$ 14.29 ^{a-P} < 0.001	381.5 $\pm$ 13.94 ^{a-P} = 0.039, b-P < 0.001

Notes: Body temperature was calculated on day-4 post infection. Data represent mean $\pm$ S.E.M for cytokine level (n = 10). a = Significantly different comparison with infected control; b = Significantly different comparison with CQ-treated; n.d. = No significantly different comparison with control groups; b.w. = body weight; CQ = chloroquine; IFN = Interferon, IL = interleukin; TNF = Tumour Necrosis Factor.

DISCUSSION

Caffeine is known for its function as a stimulant for the central nervous system in increasing focus and alertness and elevating fatigue (Barcelos *et al.* 2020). Caffeine is commonly taken in the form of drinks such as coffee or tea (Pohanka 2015). In a previous report, caffeine demonstrated its ability as an anti-plasmodial activity through an *in vitro* study where a high inhibition percentage of caffeine pyrazolines was identified against *P. falciparum* (Insuasty *et al.* 2015). It is also worth mentioning that caffeine has several properties that have been reported. This includes anti-inflammatory, antioxidant and neuroprotective, especially for the blood-brain barrier, particularly for neurodegenerative diseases (Azam *et al.* 2003; Barcelos *et al.* 2020; Yacoubi *et al.* 2000; Kolahdouzan & Hamadeh 2017).

The animal model used in this study mimics the injury of cerebral malaria. The similarity of the neurological conditions of neurodegenerative diseases with cerebral malaria, such as brain ischemia-like syndrome, excitotoxicity and inflammation, suggests that treatment with caffeine in cerebral malaria infection may also be associated with the A2AR receptor (Kolahdouzan & Hamadeh 2017). Previous studies reported that caffeine protects against neurodegenerative disease by binding to the A2AR pathway and acting as an antagonist to A2AR receptors (Chen et al. 2010). Yacoubi et al. (2000) also strengthened the fact that A2AR participation caused by the stimulant effects of caffeine at 6.25 mg/kg b.w. in the locomotor stimulatory effect in CD1 mice (Yacoubi et al. 2000). The study also stated that when the concentration of caffeine increases, the inhibition of the A2AR receptor decreases (Yacoubi et al. 2000).

Recently, plants have been used as immunomodulatory agents where natural resources have been proven to be safe and effective alternatives (Alhazmi et al. 2021). Immunomodulatory strategies are considered adjunctive therapy when the elimination of parasites cannot be achieved through antiparasitic approaches to reduce disease severity and burden (Hassan et al. 2022). Caffeine is thus a potential molecule from plant-based immunomodulators to address parasite infection as an adjunctive therapy (Kovács et al. 2021). Our findings reveal that caffeine administration into *P. berghei* ANKA-infected ICR mice effectively suppressed parasitaemia growth in red blood cells and improved the survivability of infected mice, providing proof that caffeine has good anti-malarial properties. Caffeine administration at a low dose (5 mg/kg b.w.) resulted in a significant reduction ($p < 0.05$) of parasite percentage, a significant increase in chemosuppressive percentage, and a significantly prolonged survivability of malaria-infected mice.

The result also suggested that as the concentration of caffeine increases, the chemosuppressive percentage and the survivability of infected mice decrease. Similar findings were identified from a previous study using *P. berghei* ANKA in ICR mice but with a different compound, quercetin (Ali et al. 2021). The chemosuppression percentage of quercetin at 15, 25, and 50 mg/kg b.w. was recorded as 36.1 ± 5.7 , 24.9 ± 3.8 , and 24.3 ± 1.8 %, respectively (Ali et al. 2021). Another study using andrographolide treatment in *P. berghei* NK65-infected ICR mice also reported similar findings, where at 5 mg/kg, 15 mg/kg, and 45 mg/kg b.w. showed chemosuppression at 60.17 ± 2.12 %, 58.96 ± 4.75 %, and 43.18 ± 6.28 %, respectively (Hassan et al. 2019). Our result was in line with the above-mentioned studies, which showed caffeine at 5 mg/kg, 10 mg/kg, and 20 mg/kg b.w. exhibited chemosuppression percentage of 64.18 ± 9.17 , 39.93 ± 5.63 , and 21.86 ± 5.57 %, respectively. Similar trends are also identified for the survival time as the concentration of caffeine (5 mg/kg, 10 mg/

kg and 20 mg/kg b.w.) increases, and the mean survival times of mice decrease (17, 15, and 11 days). Quercetin treatment at 15 mg/kg, 25 mg/kg and 50 mg/kg b.w. resulted at 10, 9 and 9 days and andrographolide at 5 mg/kg, 15 mg/kg and 45 mg/kg b.w. recorded mean survival times as 19, 17 and 14 days in malaria-infected mice (Ali *et al.* 2021; Hassan *et al.* 2019).

Caffeine has anti-obesity properties where caffeine can undergo cellular energy metabolism and has an antagonist effect towards adenosine receptors, which results in the inhibition of adenylyl cyclase enzyme and upregulation of cyclic adenosine monophosphate (cAMP) levels (A2AR pathway) (de Souza *et al.* 2021). Free fatty acids from the lipolysis of the cycle are then used by *Plasmodium spp.* led increases of parasite growth (Maier & van Ooij 2022). This is because *Plasmodium spp.* depends on the host to produce fatty acids (Jayabalasingham *et al.* 2010). However, the loss of body weight and body temperature were insignificant in all treatment groups compared to infected mice, as reported in the cerebral malaria murine model study (Ali *et al.* 2022; Basir *et al.* 2012).

A previous study identified that caffeine treatment at 5 mg/kg, 10 mg/kg and 20 mg/kg b.w. given together with olive oil, a good chemosuppressive percentage exhibited 79.09%, 83.64% and 87.27%, respectively (Fadare *et al.* 2024). Our study revealed lower chemosuppressive results than the previous study (Fadare *et al.* 2024). Due to the pathway of drug delivery via oral gavage and the bypass lymphatic pathway, it is possible that the drug delivery method and dilution with distilled water are the reasons for the contradiction with our findings (Yan *et al.* 2023). The study elucidated absorption using a lipid-based drug delivery system, which bypasses the first metabolic pathway into the lymphatic system (Yan *et al.* 2023). In addition, the study also stated that different dose formulations have different absorption rates, where absorption of caffeine dissolved in distilled water is slower than caffeine dissolved in olive oil (Fadare *et al.* 2024).

In addition, caffeine (5 mg/kg b.w.) reduced pro-inflammatory cytokines (TNF- α , IFN- γ , IL-18) and anti-inflammatory cytokines (IL-4 and IL-10) levels as compared to malaria-infected mice. Our results exhibited a significant reduction ($p < 0.05$) of pro-inflammatory cytokine levels (TNF- α and IFN- γ), corroborating previous findings with quercetin and andrographolide (Ali *et al.* 2021; Hassan *et al.* 2019). However, anti-inflammatory cytokine levels were significantly lower than those of infected mice for IL-10, and there was no significant difference in IL-4 levels in caffeine-treated malaria mice. Furthermore, IL-4 revealed a reduction when given treatment with CQ compared to infected mice, but it is not significant, as in a previous study (Ali *et al.* 2021). Another study by Sudi *et al.* (2018) also indicated no significant reduction of IL-4 cytokine levels when given CQ treatment.

Findings from previous studies have revealed the possible role played by each cytokine in the pathogenesis of malaria (Bhatt et al. 2021; Angulo & Fresno, 2002; Suhaini et al. 2015; Wong et al. 2015). Caffeine has also been reported to reduce cytokine production (IFN- γ , TNF- α , IL-10 and IL-18) levels in *in vitro* studies (Pohanka, 2015). A previous study of virulent *Listeria monocytogenes* in Swiss mice showed that caffeine did not exhibit any changes in IL-10 plasma level after treatment via IP injection. The study explains that the cause could be the route of administration (De Alcântara Almeida et al. 2021).

As caffeine has a biphasic effect, treatment with caffeine would not reduce pro-inflammatory cytokine levels as low as CQ treatment. Pro-inflammatory cytokine levels of caffeine treatment (TNF- α , IFN- γ , IL-18) decreased slightly (1.07-, 1.11- and 1.12-fold) as compared to infected control mice. Therefore, during malarial infection, the concentration increases from 10 mg/kg b.w. to 20 mg/kg b.w. of caffeine, resulting in excessive pro-inflammatory cytokine levels, which results in greater tissue damage with malarial presence. Thus, this causes organs to be inflamed, resulting in chronic infection. This is proven by Otha et al. (2007), who reported that 10 mg/kg b.w. of caffeine showed significant liver damage and 20 mg/kg b.w. showed a higher reading of liver damage in the acute liver study. However, caffeine at 100 mg/kg b.w. did not show any liver damage in the acute liver study (Ohta et al. 2007). The investigation of the model of ischemia-reperfusion (IR) by inhibiting A2AR shows the reduction of pERK activation and glutamate protein levels and downstream inflammatory responses (Mohamed et al. 2016). The levels of pro-inflammatory biomarkers such as NF-K β , TNF- α , IL-6, and prostaglandin E2 were lowered, while the anti-inflammatory biomarker IL-10 was increased, similar to our control group (Mohamed et al. 2016).

The A2A adenosine receptor (A2AR) activates with the binding of adenosine to the receptor, triggering a pro-inflammatory signalling pathway (Kolahdouzan & Hamadeh 2017). Once activated, A2AR triggers Gs-coupled signalling that increases cAMP and activates protein kinase A (PKA), which enhances calcium influx and glutamate release, leading to activation of the NF-K β pathway and production of inflammatory cytokines such as TNF- α and IL-6 (Kolahdouzan & Hamadeh 2017). Caffeine acts as a competitive antagonist at A2AR, blocking adenosine from binding (Kolahdouzan & Hamadeh 2017). This prevents the rise in cAMP and PKA activity, reduces calcium and glutamate-driven inflammation, and inhibits NF-K β activation. Ultimately, the cascades lowered the release of inflammatory cytokines and produced an overall anti-inflammatory and neuroprotective effect (Kolahdouzan & Hamadeh 2017).

Concentration treatments that are selected in the experiment are low-to-moderate doses of caffeine (5 mg/kg b.w, 10mg/kg b.w and 20 mg/kg b.w) and are widely used in rodent models evaluating neuroprotective, anti-inflammatory, and infection-related responses. Higher concentration of caffeine at 30 mg/kg b.w. is not chosen because there is excessive systemic toxicity (Chen *et al.* 2010).

The result shows that caffeine exhibits a dose-dependent paradox in which increasing concentration can ultimately decrease its inhibitory efficacy. At the molecular level, caffeine acts as a competitive antagonist at adenosine receptors, particularly at A₂A receptor (A₂AR) in the brain, where caffeine competes with adenosine for the same binding site (Chen *et al.* 2020; Borea *et al.* 2018). At low or acute doses, caffeine effectively binds to A₂AR without activating it, thereby blocking adenosine and pro-inflammatory signalling pathways (Chen *et al.* 2020). Thus, the blockade suppresses downstream NF- κ B activation through cAMP–PKA signalling modulation, resulting in lower inflammatory cytokine output (Liu *et al.* 2024). In this context, caffeine produces high inhibitory efficacy and reduced inflammatory signalling (Barcelos *et al.* 2020).

However, under chronic or high-dose exposure, the brain and peripheral tissues activate homeostatic feedback mechanisms. Prolonged A₂AR blockade by caffeine induces adaptive upregulation of adenosine receptors, increasing A₂AR density and thereby reducing caffeine's relative blocking efficiency (de Souza Gonçalves *et al.* 2017; Eichwald *et al.* 2023). With more A₂AR available, caffeine must compete against higher adenosine levels for a larger receptor pool, diminishing its antagonistic efficacy and contributing to pharmacological tolerance (Chen *et al.* 2020). As A₂AR signalling regains influence, NF- κ B-mediated inflammatory pathways can become reactivated, promoting increased cytokine production compared with the low-concentration (Liu *et al.*, 2024; Marcinek *et al.* 2024). Thus, these may lead to higher production of inflammatory cytokines.

Taken together, findings demonstrate that the effective dose of caffeine can suppress parasite growth and modulate the cytokine levels in malaria-infected ICR mice. The role of caffeine in enhancing the immune response against *P. berghei* ANKA, particularly its effects on pro-inflammatory cytokine production, demonstrated that caffeine interactions with cells are an area for further exploration. Additionally, the study raises the significance of investigating the pharmacodynamics and potential mechanisms of caffeine's action against malarial parasites. Theoretically, caffeine at 5 mg/kg b.w. can inhibit A₂AR, which reduces the production of inflammatory cytokines and slows down malarial infection growth.

This study acknowledges several limitations. Histopathological analyses of the brain, liver, and spleen were not performed to compare infected tissues with treated animals. Such analyses would have provided stronger evidence regarding the potential of caffeine to improve cerebral malaria conditions. Additionally, oxidative stress biomarkers, including malondialdehyde (MDA) and reduced glutathione (GSH), were not evaluated. Another limitation is the use of only male mice. Female mice are not used due to the occurrence of hormonal imbalances, such as estrogen, which increases oxidative stress, influences malarial parasite growth and leads to cytokine storm, potentially affecting the consistency of results within the experimental groups (Wu et al. 2025).

Further investigation into the pharmacodynamic profile of caffeine in the context of malaria treatment is also recommended to better understand its mechanism of action and therapeutic potential. Future studies should explore the potential of combination therapies involving caffeine and standard antimalarial drugs, such as artesunate or CQ, to evaluate possible synergistic effects.

CONCLUSION

The study has shown that caffeine possesses anti-malarial properties and is able to modulate inflammatory cytokine levels. Treatment with caffeine at 5 mg/kg b.w. in *P. berghei* ANKA-infected ICR mice was able to slow down parasite growth as compared to 10 mg/kg b.w. and 20 mg/kg b.w. of caffeine treatment. Notably, the low-dose caffeine exhibited a good chemosuppressive effect (> 60%) and was able to significantly prolong the survival time of malaria-infected ICR mice (17 days). Furthermore, caffeine at 5 mg/kg b.w. modulated cytokine levels of the pro-inflammatory (TNF- α , IFN- γ and IL-18) and anti-inflammatory (IL-4 and IL-10) cytokines. This results in the anti-malarial and cytokine-modulating properties in ICR mice. The author also plans to identify the mechanism of caffeine, i.e., whether caffeine can block the A2AR pathway.

AUTHORS' CONTRIBUTIONS

Ahmad Zakwan Noriman: Writing – original draft, review and editing, conceptualisation, formal analysis, methodology, investigation.

Mohd Ridzuan Mohd Abd Razak: Resources, supervision, writing – review.

Wan Nurhayati Wan Hanafi: Supervision, writing – review.

Rusliza Basir: Resources, supervision, writing – review.

Hani Kartini Agustar: Supervision, writing – review.

Wan Rozianoor Mohd Hassan: Writing – review and editing, conceptualisation, validation, supervision and funding.

AUTHOR(S)' DISCLOSURE (CONFLICT OF INTEREST) STATEMENT(S)

The authors declare no potential conflicts of interest.

FUNDING STATEMENT

This research was funded by the Ministry of Higher Education, Malaysia (FRGS/1/2021/STG01/UiTM/02/5) and Journal Support Fund, Universiti Teknologi MARA, Shah Alam, Malaysia.

REFERENCES

- Akinyemi D O, Olorunfemi A B, Akinola O, Abiodun O O, Afolabi O K, Thomas B N, Gbotosho G O and Ojurongbe O. (2025). Antimalarial, antioxidant and hepatoprotective activities of ethanol extract of *Blighia sapida* K.D. Koenig (*Sapindaceae*) in *Plasmodium berghei*-infected mice. *Acta Parasitologica* 70: 1–21. <https://doi.org/10.1007/s11686-025-01162-4>
- Alhazmi H A, Najmi A, Javed S A, Sultana S, Bratty M A, Makeen H A, Meraya A M, Ahsan W, Mohan S, Taha M M E and Khalid A. (2021). Medicinal plants and isolated molecules demonstrating immunomodulation activity as potential alternative therapies for viral diseases including COVID-19. *Frontiers in Immunology* 12: 1–24. <https://doi.org/10.3389/fimmu.2021.637553>
- Ali A H, Hassan W R M, Dahari D E, Embi N, Sidek H M, Basir R, Agustar H K and Latip J. (2022). Pencirian jangkitan *Plasmodium berghei* nk65 terhadap mencit ICR sebagai model jangkitan malaria teruk. *Sains Malaysiana* 51(10): 3481–3493. <https://doi.org/10.17576/jsm-2022-5110-29>
- Ali A H, Sudi S, Shi-Jing N, Hassan W R M, Basir R, Agustar H K, Embi N, Sidek H M and Latip J. (2021). Dual anti-malarial and GSK3B-mediated cytokine-modulating activities of quercetin are requisite of its potential as a plant-derived therapeutic in malaria. *Pharmaceuticals* 14(3): 1–15. <https://doi.org/10.3390/ph14030248>
- Almeida I D A, Dorvigny B M, Tavares L S, Santana L N and Lima-Filho J V. (2021). Anti-inflammatory activity of caffeine (1,3,7-trimethylxanthine) after experimental challenge with virulent *Listeria monocytogenes* in Swiss mice. *International Immunopharmacology* 100: 1–7. <https://doi.org/10.1016/j.intimp.2021.108090>
- Angulo I and Fresno M. (2002). Cytokines in the pathogenesis of and protection against malaria. *Clinical and Vaccine Immunology* 9(6): 1145–1152. <https://doi.org/10.1128/cdli.9.6.1145-1152.2002>

- Azam S, Hadi N, Khan N U and Hadi S M. (2003). Antioxidant and prooxidant properties of caffeine, theobromine and xanthine. *Medical Science Monitor* 9(9): 325–330.
- Barcelos R P, Lima F D, Carvalho N R, Bresciani G and Royes L F. (2020). Caffeine effects on systemic metabolism, oxidative-inflammatory pathways, and exercise performance. *Nutrition Research* 80: 1–17. <https://doi.org/10.1016/j.nutres.2020.05.005>
- Basir R, Rahiman S, Hasballah K, Chong W, Talib H, Yam M, Jabbarzare M, Tie T, Othman F, Moklas M, Abdullah W and Ahmad Z. (2012). *Plasmodium berghei* ANKA infection in ICR mice as a model of cerebral malaria. *Iranian Journal of Parasitology* 7(4): 62–74.
- Bhatt D, Kumar S, Kumar P, Bisht S, Kumar A, Maurya A K, Pal A and Bawankule D U. (2022). Rutin ameliorates malaria pathogenesis by modulating inflammatory mechanism: an *in vitro* and *in vivo* study. *Inflammopharmacology* 30(1): 159–171. <https://doi.org/10.1007/s10787-021-00920-w>
- Borea P A, Gessi S, Merighi S, Vincenzi F and Varani K. (2018). Pharmacology of adenosine receptors: The state of the art. *Physiological Reviews* 98(3): 1591–1625. <https://doi.org/10.1152/physrev.00049.2017>
- Chen J and Schwarzschild M A. (2020). Do caffeine and more selective adenosine A2A receptor antagonists protect against dopaminergic neurodegeneration in Parkinson's disease? *Parkinsonism & Related Disorders* 80: 45–53. <https://doi.org/10.1016/j.parkreldis.2020.10.024>
- Chen X, Ghribi O and Geiger J D. (2010). Caffeine protects against disruptions of the blood-brain barrier in animal models of Alzheimer's and Parkinson's diseases. *Journal of Alzheimer's Disease* 20(s1): 127–141. <https://doi.org/10.3233/jad-2010-1376>
- Craig A G, Grau G E, Janse C, Kazura J W, Milner D, Barnwell J W, Turner G and Langhorne J. (2012). The role of animal models for research on severe malaria. *PLoS Pathogens* 8(2):1–10. <https://doi.org/10.1371/journal.ppat.1002401>
- Das R G and North D. (2007). Implications of experimental technique for analysis and interpretation of data from animal experiments: outliers and increased variability resulting from failure of intraperitoneal injection procedures. *Laboratory Animals* 41(3): 312–320. <https://doi.org/10.1258/002367707781282802>
- De Souza Gonçalves L S, Coelho D B, de Oliveira G V and Rodrigues L O C. (2017). Dispelling the myth that habitual caffeine consumption increases tolerance to the ergogenic effects of caffeine. *Journal of Applied Physiology* 123(1): 213–220. <https://doi.org/10.1152/jappphysiol.01029.2016>
- De Souza F B, Novaes R D, Santos C F F, De Deus F A, Santos F C, Ladeira L C M, Gonçalves R V, Bastos D S S, Souza A C F, Machado-Neves M and Santos E C D. (2021). High-fat diet and caffeine interact to modulate bone microstructure and biomechanics in mice. *Life Sciences* 276: 1–10. <https://doi.org/10.1016/j.lfs.2021.119450>
- Dorvigny B M, Tavares L S, De Almeida I A, Santana L N, Silva E D S, De Souza J K U, Soares A F, Da Silva Júnior V A and Lima-Filho J V. (2021). Antiinflammatory and antiinfective effect of caffeine in a mouse model of disseminated salmonellosis. *Phytotherapy Research* 36(4): 1652–1663. <https://doi.org/10.1002/ptr.7349>
- Eichwald T, Solano A F, Souza J, De Miranda T B, Carvalho L B, Sanna P L D S, Da Silva R A F and Latini A. (2023). Anti-inflammatory effect of caffeine on muscle under lipopolysaccharide-induced inflammation. *Antioxidants* 12(3): 1–54. <https://doi.org/10.3390/antiox12030554>

- Fadare O, Omisore N, Fadare R, Oduwale A, Awofisayo O, Ogundolie F, Salaria D, Rolta R and Adesanwo J (2024). Antimalarial activity of caffeine, orally administered with a lipid-based formulation in a Murine model. *Ife Journal of Science* 26(2): 277–291. <https://doi.org/10.4314/ij.s.v26i2.5>
- Habibi P, Shi Y, Grossi-De-Sa M F and Khan, I. (2022). Plants as sources of natural and recombinant antimalaria agents. *Molecular Biotechnology* 64(11): 1177–1197. <https://doi.org/10.1007/s12033-022-00499-9>
- Hassan W R M, Ali A H, Basir R, Embi N and Sidek H M. (2022). Medicinal plants with antimalarial activities mediated via glycogen synthase kinase-3 beta (GSK3B) inhibition. *Tropical Biomedicine*, 39(3): 384–393. <https://doi.org/10.47665/tb.39.3.008>
- Hassan W R M, Basir R, Ali A H, Embi N and Sidek H M. (2019). Anti-malarial and cytokine-modulating effects of andrographolide in a murine model of malarial infection. *Tropical Biomedicine* 36(3): 776–791.
- Huang B, Pearman E, and Kim C. (2015). Mouse models of uncomplicated and fatal malaria. *BIO-PROTOCOL* 5(13): 1–12. <https://doi.org/10.21769/bioprotoc.1514>
- Insuasty B, Ramírez J, Becerra D, Echeverry C, Quiroga J, Abonia R, Robledo S M, Vélez I D, Upegui Y, Muñoz J A. et al. (2015). An efficient synthesis of new caffeine-based chalcones, pyrazolines and pyrazolo[3,4-b][1,4]diazepines as potential antimalarial, antitrypanosomal and antileishmanial agents. *European Journal of Medicinal Chemistry* 93: 401–413. <https://doi.org/10.1016/j.ejmech.2015.02.040>
- Jayabalasingham B, Menard R and Fidock D A. (2010). Recent insights into fatty acid acquisition and metabolism in malarial parasites. *F1000 Biology Reports*, 2: 1–12. <https://doi.org/10.3410/b2-24>
- Kolahdouzan M and Hamadeh M J. (2017). The neuroprotective effects of caffeine in neurodegenerative diseases. *CNS Neuroscience & Therapeutics* 23(4): 272–290. <https://doi.org/10.1111/cns.12684>
- Kovács E G, Alatshan A, Budai M M, Czimmerer Z, Bíró E and Benkő S. (2021). Caffeine has different immunomodulatory effect on the cytokine expression and NLRP3 inflammasome function in various human macrophage subpopulations. *Nutrients* 13(7): 1–20. <https://doi.org/10.3390/nu13072409>
- Kwiatkowski D and Bate C. (1995). Inhibition of tumour necrosis factor (TNF) production by antimalarial drugs used in cerebral malaria. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 89(2): 215–216. [https://doi.org/10.1016/0035-9203\(95\)90501-4](https://doi.org/10.1016/0035-9203(95)90501-4)
- Lee W, Cheong F W, Amir A, Lai M Y, Tan J H, Phang W K, Shahari S and Lau Y. (2022). *Plasmodium knowlesi*: the game changer for malaria eradication. *Malaria Journal* 21(1): 1–20. <https://doi.org/10.1186/s12936-022-04131-8>
- Leitner W W, Bergmann-Leitner E S and Angov E. (2010). Comparison of *Plasmodium berghei* challenge models for the evaluation of pre-erythrocytic malaria vaccines and their effect on perceived vaccine efficacy. *Malaria Journal* 9(1): 1–12. <https://doi.org/10.1186/1475-2875-9-145>
- Liu W Li Q, Fang W, Cai L, Wang Z, Kou B, Zhou C, Zhou Y, Yao Z, Wei M and Zhang, S. (2024). A2AR regulate inflammation through PKA/NF-KB signaling pathways in intervertebral disc degeneration. *European Journal of Medical Research* 29(1): 1–13. <https://doi.org/10.1186/s40001-024-02028-7>
- Maier A G and Ooij, C V. (2022). The role of cholesterol in invasion and growth of malaria parasites. *Frontiers in Cellular and Infection Microbiology* 12: 1–14. <https://doi.org/10.3389/fcimb.2022.984049>
- Marcinek K, Luzak B, and Rozalski M. (2024). The effects of caffeine on blood platelets and the cardiovascular system through adenosine receptors. *International Journal of Molecular Sciences* 25(16): 1–17. <https://doi.org/10.3390/ijms25168905>

- Mekonnen L B. (2015). *In vivo* antimalarial activity of the crude root and fruit extracts of *Croton macrostachyus* (Euphorbiaceae) against *Plasmodium berghei* in mice. *Journal of Traditional and Complementary Medicine* 5(3): 168–173. <https://doi.org/10.1016/j.jtcme.2014.07.002>
- Mohamed R, Agha A, Abdel-Rahman A, and Nassar N. (2015). Role of adenosine A2A receptor in cerebral ischemia reperfusion injury: Signaling to phosphorylated extracellular signal-regulated protein kinase (pERK1/2). *Neuroscience* 314: 145–159. <https://doi.org/10.1016/j.neuroscience.2015.11.059>
- Ogbeide O K, Sajere O, Obukuru I S, Akedayi P, Omofe N O, Unachukwu C I, Mbonu R, Aghedo N O, Iyekowa O and Uadia J. (2024). Comparative studies on the bioactive chemical composition, antioxidant and antimalarial potentials of *Picralima nitida* leaf and stem bark extracts. *Benin Journal of Physical Sciences* 1(1): 231–250.
- Ohta A, Lukashev D, Jackson E K, Fredholm B B, and Sitkovsky M. (2007). 1,3,7-Trimethylxanthine (caffeine) may exacerbate acute inflammatory liver injury by weakening the physiological immunosuppressive mechanism. *The Journal of Immunology* 179(11): 7431–7438. <https://doi.org/10.4049/jimmunol.179.11.7431>
- Parija S, and Antony H. (2016). Antimalarial drug resistance: An overview. *Tropical Parasitology* 6(1): 30–41. <https://doi.org/10.4103/2229-5070.175081>
- Percie du Sert N, Hurst V, Ahluwalia A, Alam S, Avey M T, Baker M, Browne W J, Clark A, Cuthill I C, Dirnagl U, et al. (2020). The ARRIVE guidelines 2.0: Updated guidelines for reporting animal research. *PLoS Biology* 18(7): 1–12. <https://doi.org/10.1371/journal.pbio.3000410>
- Peters W and Robinson B L. (1992). The chemotherapy of rodent malaria. XLVII. Studies on pyronaridine and other Mannich base antimalarials. *Annals of Tropical Medicine and Parasitology* 86(5): 455–465. <https://doi.org/10.1080/00034983.1992.11812694>
- Picot S, Peyron F, Donadille A, Vuillez J P, Barbe G, and Ambroise-Thomas P. (1993). Chloroquine-induced inhibition of the production of TNF, but not of IL-6, is affected by disruption of iron metabolism. *Parasitology Today* 9(12): 443. [https://doi.org/10.1016/0169-4758\(93\)90096-x](https://doi.org/10.1016/0169-4758(93)90096-x)
- Pohanka M. (2014). Caffeine downregulates antibody production in a mouse model. *Journal of Applied Biomedicine* 13(1): 1–6. <https://doi.org/10.1016/j.jab.2014.09.001>
- Rasmussen C, Alonso P and Ringwald P. (2021). Current and emerging strategies to combat antimalarial resistance. *Expert Review of Anti-infective Therapy* 20(3): 353–372. <https://doi.org/10.1080/14787210.2021.1962291>
- Sonsalla P K, Wong L, Harris S L, Richardson J R, Khobahy I, Li W, Gadad B S and German D C. (2012). Delayed caffeine treatment prevents nigral dopamine neuron loss in a progressive rat model of Parkinson's disease. *Experimental Neurology* 234(2): 482–487. <https://doi.org/10.1016/j.expneurol.2012.01.022>
- Sudi S, Hassan W R M, Ali A H, Basir R, Embi N and Sidek H M. (2018). A derivative of cinnamic acid, methyl-4-hydroxycinnamate modulates inflammatory cytokine levels in malaria-infected mice through inhibition of GSK3B. *Malaysian Applied Biology* 47(3): 153–157.
- Suhaini S, Liew S Z, Norhaniza J, Lee P C, Jualang G, Embi N and Hasidah, M. S. (2015). Anti-malarial and anti-inflammatory effects of *Gleichenia truncata* mediated through inhibition of GSK3B. *Tropical Biomedicine* 32(3): 419–433.
- Temple J L, Bernard C, Lipshultz S E, Czachor J D, Westphal J A and Mestre M A. (2017). The safety of ingested caffeine: A comprehensive review. *Frontiers in Psychiatry* 8: 1–19. <https://doi.org/10.3389/fpsy.2017.00080>

- Titanji V P, Zofou D and Ngemenya M N. (2008). The antimalarial potential of medicinal plants used for the treatment of malaria in Cameroonian folk medicine. *African Journal of Traditional, Complementary, and Alternative Medicines* 5(3): 302–321.
- Vaou N, Stavropoulou E, Voidarou C, Tsigalou C and Bezirtzoglou E. (2021). Towards advances in medicinal plant antimicrobial activity: A review study on challenges and future perspectives. *Microorganisms* 9(10): 20–41. <https://doi.org/10.3390/microorganisms9102041>
- Verhave P, Eenige, R V & Tiebosch I. (2024). Methods for applying blinding and randomisation in animal experiments. *Laboratory Animals* 58(5): 419–426. <https://doi.org/10.1177/00236772241272991>
- Wong S K, Jann M L S, Sudi S, Hassan W R M, Chin L P, Embi N, Sidek H M. (2015). Anti-malarial and anti-inflammatory effects of *Gynura procumbens* are mediated by kaempferol via inhibition of glycogen synthase kinase-3 β (GSK3B). *Sains Malaysiana* 44(10): 1489–1500. <https://doi.org/10.17576/jsm-2015-4410-15>
- WHO (2024). *World Malaria Report 2024: Addressing inequity in the global Malaria response*. <https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2024>
- Wu Y, Chen Y, Liu F, and Li K. (2025). The immunomodulatory role of estrogen in malaria: A review of sex differences and therapeutic implications. *Immunity Inflammation and Disease* 13(2): 1–11. <https://doi.org/10.1002/iid3.70148>
- Yacoubi M E, Ledent C, Ménard J, Parmentier M, Costentin J and Vaugeois, J. (2000). The stimulant effects of caffeine on locomotor behaviour in mice are mediated through its blockade of adenosine A2A receptors. *British Journal of Pharmacology* 129(7): 1465–1473. <https://doi.org/10.1038/sj.bjp.070.3170>
- Yan S, Cheng Y, Li L, Zhong C, Chen C and Gao X. (2023). Lipid-based formulations: A promising approach for poorly soluble drug delivery via the intestinal lymphatic system. *Journal of Drug Delivery Science and Technology* 87: 104770. <https://doi.org/10.1016/j.jddst.2023.104770>
- Zhou W, Wang H, Yang Y, Chen Z, Zou C and Zhang, J. (2020). Chloroquine against malaria, cancers and viral diseases. *Drug Discovery Today* 25(11): 2012–2022. <https://doi.org/10.1016/j.drudis.2020.09.010>

APPENDIX

Appendix A: Physical signs throughout 30 days monitoring

Physical signs	Category	Days appeared					
		Normal	Infected mice (negative control)	CQ-treated (10 mg / kg b.w.) (positive control)	Caffeine-treated		
					5 mg/kg b.w.	10 mg/kg b.w.	20 mg/kg b.w.
Lethargic	Absent	Day 30	Day 2	Day 30	Day 8	Day 3	Day 3
	Mild	0	Day 3	0	Day 9	Day 4	Day 4
	Moderate	0	Day 4	0	Day 11	Day 5	Day 5
	Severe	0	Day 5	0	Day 15	Day 8	Day 6
Piloerection	Absent	Day 30	Day 2	Day 30	Day 5	Day 2	Day 2
	Mild	0	Day 3	0	Day 6	Day 3	Day 3
	Moderate	0	Day 4	0	Day 8	Day 4	Day 4
	Severe	0	Day 5	0	Day 12	Day 5	Day 5
Reduced locomotor activity	Absent	Day 30	Day 2	Day 30	Day 6	Day 3	Day 2
	Mild	0	Day 3	0	Day 7	Day 4	Day 3
	Moderate	0	Day 4	0	Day 9	Day 5	Day 4
	Severe	0	Day 5	0	Day 11	Day 5	Day 5
Urine colour	Absent	Day 30	Day 2	Day 30	Day 8	Day 4	Day 3
	Mild	0	Day 3	0	Day 9	Day 5	Day 4
	Moderate	0	Day 4	0	Day 11	Day 6	Day 5
	Severe	0	Day 5	0	Day 14	Day 9	Day 7

Notes: Physical signs changes were observed throughout 30 days. The data represent the days appearance on mice. CQ = chloroquine.