



Length-Weight Relationship, Size Distribution and Sexual Maturity of Climbing perch, *Anabas testudineus* (Bloch, 1792) From Different Habitats in Malaysia

Authors:

Mohamad Jalilah*, Victor Tosin Okomoda, Khor Waiho, Hidayah Manan, Hassan Anuar, Thumronk Amornsakun and Musa Nadirah*

***Correspondence:** nadirah@umt.edu.my; jalilah@umt.edu.my

Submitted: 12 February 2025; **Accepted:** 21 October 2025;
Published: 31 March 2026

To cite this article: Jalilah Mohamad, Victor Tosin Okomoda, Khor Waiho, Hidayah Manan, Hassan Anuar, Thumronk Amornsakun and Nadirah Musa. (2026). Length-weight relationship, size distribution and sexual maturity of climbing perch, *Anabas testudineus* (Bloch, 1792) from different habitats in Malaysia. *Tropical Life Sciences Research* 37(1): 157–184. <https://doi.org/10.21315/tlsr2026.37.1.8>

To link to this article: <https://doi.org/10.21315/tlsr2026.37.1.8>

HIGHLIGHTS

- Growth and reproductive performance of climbing perch were influenced by habitat type and seasonal variations.
- Fish in irrigation canals during the rainy season exhibited the largest size and enhanced gonadal development.
- High dissolved oxygen and pH were correlated with larger fish, while rising temperatures were linked to smaller sizes.
- Findings underscore the importance of habitat quality and seasonal changes in shaping wild climbing perch populations.



Length-Weight Relationship, Size Distribution and Sexual Maturity of Climbing Perch, *Anabas testudineus* (Bloch, 1792) From Different Habitats in Malaysia

¹Mohamad Jalilah*, ^{1,2,3}Victor Tosin Okomoda, ^{1,4,6,7}Khor Waiho, ¹Hidayah Manan, ⁵Hassan Anuar, ⁸Thumronk Amornsakun and ⁵Musa Nadirah*

¹Higher Institution Center of Excellence (HICoE), Institute of Tropical Aquaculture and Fisheries, Universiti Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia

²WorldFish, Ibadan, Nigeria. c/o International Institute of Tropical Agriculture (IITA), PMB 5320, Oyo Road Ibadan 200001, Oyo State, Nigeria

³Department of Fisheries and Aquaculture, College of Forestry and Fisheries, Joseph Sarwuan Tarka University (formerly Federal University of Agriculture Makurdi), Makurdi P.M.B. 2373 Makurdi, Nigeria

⁴STU-UMT Joint Shellfish Research Laboratory, Shantou University, Shantou, Guangdong, China

⁵Faculty of Fisheries and Food Science, Universiti Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia

⁶Centre for Chemical Biology, Universiti Sains Malaysia, 11800 Pulau Pinang, Malaysia

⁷Department of Aquaculture, Faculty of Fisheries, Kasetsart University, Bangkok, Thailand

⁸Faculty of Science and Technology, Prince of Songkhla University, Pattani campus, Amphure Muang, 94000 Pattani, Thailand

Submitted: 12 February 2025; **Accepted:** 21 October 2025; **Published:** 31 March 2026

To cite this article: Jalilah Mohamad, Victor Tosin Okomoda, Khor Waiho, Hidayah Manan, Hassan Anuar, Thumronk Amornsakun and Nadirah Musa. (2026). Length-weight relationship, size distribution and sexual maturity of climbing perch, *Anabas testudineus* (Bloch, 1792) from different habitats in Malaysia. *Tropical Life Sciences Research* 37(1): 157–184. <https://doi.org/10.21315/tlsr2026.37.1.8>

To link to this article: <https://doi.org/10.21315/tlsr2026.37.1.8>

Abstract: The growth and reproductive performance of fish are influenced by ecological factors. This study investigated the biology of the climbing perch, *Anabas testudineus*, in relation to habitat. A total of 2,880 climbing perch were collected from three habitats (irrigation canal, swamp and paddy field) in Peninsular Malaysia during the dry and rainy seasons, and their biometric measurements (total length and body weight) and sexual maturity indices were measured. Two-way Classification Analysis (TCA) revealed significant sexual dimorphism in all morphometric parameters, significant habitat effects on fish size, and pronounced seasonal variation in environmental parameters, while condition factor and seasonal effects on morphometrics were not significant. Results showed that body size was largest in fish from irrigation canals during the rainy season, with males averaging 12.84 ± 1.17 cm in total length (32.38 ± 4.4 g body weight) and females averaging 13.70 ± 1.83 cm (52.84 ± 12.54 g), consistent with the upper size range reported for this species. Fish from irrigation canals also exhibited more advanced gonadal development, confirmed by higher gonadosomatic indices (GSI) and histological evidence of vitellogenic oocytes in females and spermatozoa-filled seminiferous lobules in males. Principal Component Analysis (PCA) indicated that larger body sizes and advanced gonadal development were associated with higher dissolved oxygen and pH, whereas smaller sizes correlated with elevated water temperature. This study highlights

the critical roles of sex, habitat quality and seasonal variation in shaping the growth and reproductive traits of wild climbing perch populations and provides evidence-based insights for fisheries management in tropical freshwater ecosystems.

Keywords: Growth, Length-Weight Relationship, Condition Factor, Reproductive Performance, Size at Maturity

Abstrak: Tumbesaran dan prestasi pembiakan ikan dipengaruhi oleh faktor-faktor ekologi. Kajian ini menyelidik beberapa aspek biologi ikan puyu, *Anabas testudineus* yang berkaitan dengan habitat mereka. Sejumlah 2,880 ikan puyu telah dikumpulkan dari tiga habitat (iaitu terusan pengairan, paya dan sawah padi) di Semenanjung Malaysia semasa musim panas dan hujan, dan parameter morfometrik serta indeks kematangan seksual mereka diukur dan dianggarkan. Keputusan menunjukkan perbezaan yang signifikan ($P < 0.05$) dalam kebanyakan parameter antara ikan jantan dan betina, serta merentasi habitat dan musim. Saiz badan ikan puyu yang disampel semasa musim hujan dari terusan pengairan adalah paling besar berbanding musim panas dan habitat lain (jantan, 12.84 ± 1.17 cm panjang keseluruhan dan 32.38 ± 4.4 g berat badan; betina, 13.70 ± 1.83 cm panjang keseluruhan dan 52.84 ± 12.54 g berat badan). Ikan dari terusan pengairan semasa musim hujan menunjukkan saiz badan terbesar dan perkembangan gonad yang lebih bagus, seperti yang dibuktikan oleh indeks gonadosomatik (GSI) yang lebih tinggi dan gamet yang terbeza dengan baik berbanding dengan ikan dari habitat lain. Trend kualiti air menunjukkan bahawa saiz badan yang lebih besar dikaitkan dengan tahap oksigen terlarut (DO) dan pH yang lebih tinggi, manakala saiz badan yang lebih kecil dikaitkan dengan suhu air yang tinggi. Kajian ini menekankan peranan penting kualiti habitat, variasi bermusim, dan jantina dalam membentuk ciri-ciri tumbesaran dan pembiakan populasi ikan puyu liar.

Kata kunci: Tumbesaran, Hubungan Panjang-Berat, Faktor Keadaan, Prestasi Pembiakan, Saiz pada Kematangan

INTRODUCTION

The climbing perch, *Anabas testudineus* (Bloch, 1792) (Perciformes: Anabantoidae: Anabantidae) is a commercially important fish in Malaysia, with a market price ranging between RM18 and RM20 per kilogram (Chaturvedi et al. 2015; Marimuthu et al. 2009a; Hassan 2017; Syazwani 2020) and highly nutritious (Uttam et al. 2005; Tay et al. 2006; Mohanty et al. 2016; Paul et al. 2017; Akter et al. 2020). Besides being a protein source, the species also serves as a biological control agent for mosquito larvae, including *Aedes*, *Anopheles* and *Culex*, due to its tolerance of poor water quality (Hussain 2005; Chandra et al. 2008; Khairul 2011).

Climbing perch are found mostly in freshwater ecosystems; inhabiting habitats such as paddy fields, swamps, reservoirs, ditches, ponds, marshland, canals and irrigation canals, as well as streams that are filled with vegetation (Iwata & Kiguchi 2003; Das et al. 2009; Froese & Pauly 2019). They possess labyrinth organs enabling aerial respiration and can withstand drought by burrowing into mud (Jacob 2005; Hitchcock 2008; Morioka et al. 2009b). However, wild populations are declining due to habitat destruction caused by overfishing, urbanisation, agriculture, industry and pollution (Khatune-Jannat et al. 2012; Hossain et al. 2015).

Knowledge of their growth and reproduction is vital for developing breeding techniques and improving seed production (Kagawa 2013). Many studies have examined reproductive biology, including gonadal development (Uddin *et al.* 2016; Priyatha & Chitra 2022), fecundity (Marimuthu *et al.* 2009b), breeding behaviour (Zworykin 2012), larval morphology and growth (Amornsakun *et al.* 2005; Morioka *et al.* 2009a), culture systems (Rahman & Marimuthu 2010; Uddin *et al.* 2016; Anantharaja *et al.* 2017), feeding requirements (Hossain *et al.* 2012) and disease control (Sumon *et al.* 2015; Tran *et al.* 2024; Shahriar *et al.* 2024). Yet, there remains limited information on ecological interactions such as water quality, habitat diversity and climate influences.

The capability of a species to reproduce successfully under a rapidly changing environmental condition is crucial to ensuring its continuity in an ecosystem (Moyle & Crech 2000; Le Doux-Bloom *et al.* 2021; Morales-Gamba *et al.* 2021). This is because the life history strategies such as development, growth and reproduction depend largely on the environmental conditions of the habitat (Magalhaes *et al.* 2003; Olden *et al.* 2006). Distinct ecological variables drive adaptive strategies involving developmental, behavioural and physiological changes (Andrade & Braga 2005; Amtyaz Khan *et al.* 2013; Fafioye & Ayodele 2018; Ahmadi 2019).

A study of the length-weight relationship (LWR), size distribution and sexual maturity of climbing perch is essential to assess the growth patterns, reproductive potential and overall health of the population on this species, which plays a vital role in sustainable management as well as aquaculture practices. According to Ndobe *et al.* (2019), LWR reveals growth patterns, whether isometric (weight proportional to length) or allometric (weight grows disproportionately). This also relates to condition factor, an indicator of population health (Khatune-Jannat *et al.* 2012). LWR knowledge supports aquaculture by identifying optimal conditions for growth such as diet and environment (Hamid *et al.* 2015). Size distribution provides insight into age structure and recruitment success, while skewed distribution may reflect overfishing or habitat stress (Wilson *et al.* 2010).

Moreover, the investigation of sexual maturity is basic to estimating the reproductive capacity of the species. Because life history information such as the age and size at first maturity speaks to population resiliency, it is critical for establishing successful harvesting regulations including the minimum size limit set to avoid the take of immature individuals, thus enabling the population to be replenished (Cerdenares-Ladrón *et al.* 2011; Perez-Palafox *et al.* 2022). In aquaculture, the utilisation of climbing perch reproductive biology information facilitates control of breeding programs for maximum production and sustainability (Mawa *et al.* 2022). Sexual maturity also reflects environmental effects such as temperature

(Kuparinen *et al.* 2011), water quality (Marius *et al.* 2023) and food availability (Yoneda & Wright 2005), supporting both aquaculture and conservation.

Similarly, tracking LWR, size distribution and sexual maturity can provide insight for developing targeted management plans, such as seasonal fishing bans or the designation of protected breeding grounds (Russell *et al.* 2011). Due to the wide distribution range of climbing perch throughout Southeast Asia and its importance in both subsistence and commercial fisheries (Kumar *et al.* 2012), this is particularly relevant. Knowing these biological parameters in aquaculture is essential for effective stock management, improving breeding efficiency and making production practices sustainable and environmentally friendly (Rahman & Samat 2021). Thus, these parameters are indispensable for improving fishery sustainability and conservation of climbing perch.

Other factors such as food availability/quality, population density, predation force, physical and physiological stressors as well as disease threats also play important role in shaping a species' physiological characteristics (Khatune-Jannat *et al.* 2012; Abdel-Tawwab *et al.* 2015; Ndobe *et al.* 2019). Yet, information on climbing perch remains limited, especially in Malaysia, despite extensive studies in other Asian countries (Mukherjee *et al.* 2003; Jacob 2005; Sarkar *et al.* 2005; Atal *et al.* 2009; Morioka *et al.* 2009b; Kumar *et al.* 2013; Ahmadi 2019). Therefore, this study was designed to investigate the size distribution, LWR, and maturity of climbing perch across different seasons and habitats of Peninsular Malaysia. Findings will provide baseline data on growth, maturity and spawning, serving as reference for stock assessment, management and conservation.

MATERIALS AND METHODS

Sampling Site and Condition

A total of 2,880 adult-sized climbing perch (1:1 male-to-female ratio) were collected from four states in Malaysia. Two states from the north-eastern [i.e., Kelantan (5.2500°N, 102.0000°E) and Terengganu (4.7500°N, 103.0000°E)], and north-western Peninsular Malaysia [i.e., Kedah (6.1283° N, 100.3628°E) and Perlis (6.5000°N, 100.3217°E)] were selected for this study (Fig. 1). A total of 60 fish per state were captured by trapping (Trap size; 2 × 2 × 2 feet and mesh size of 1 inch) angling (hook size; 10–12 and worm as bait) and netting (mesh size; 1 inch) monthly over a period of one-year (both dry and rainy season). According to the Malaysian Meteorological Department (2012; 2013), the dry season in the eastern and northern regions of Peninsular Malaysia was observed from February

to July and from January to June, respectively. The rainy season, on the other hand, extended from August to January in the eastern region and from July to December in the northern region. The physico-chemical characteristic of water (water temperature, dissolved oxygen and pH) in each sampling site were recorded *in-situ* using an hand-held YSI meter (YSI model 556 MPS, USA) in three replicates.




FIGURE 1: Map of Peninsular Malaysia showing the sampling locations across the eastern and northern states. In Kelantan, the sites were (A) Tumpat, (B) Bachok and (C) Pasir Putih, while in Terengganu the sites included (D) Besut, (E) Setiu and (F) Kuala Terengganu. In the northern region, sampling was conducted in Kedah at (G) Kubang Pasu, (H) Kuala Muda and (I) Kulim, and in Perlis at (J) Beseri, (K) Padang Besar and (L) Mata Ayer.

Sampling Habitats

At the different states, fish were captured from three habitats namely, paddy fields, irrigation canals and swamps. Paddy field selected are paddy cultivation areas with a water depth of about 0.40 ± 0.05 m with soft, muddy substrate (following habitat descriptions adapted from Das 2002). The irrigation canals selected were 1.65 ± 0.09 m in water depth. They have muddy sandy substrate while the sides of the canals are dominated with water grasses (similar to the methods described by Katano *et al.* (2003) and Thompson and Larsen (2004). The swamp area selected have a static cloudy water with depth of about 1.70 ± 0.12 m. The substrates are mostly muddy, covered by woody debris (based on standard sampling habitat classification in Boyd and Tucker [1992] and Christensen *et al.* [1996]). The water depths reported reflect the natural and typical conditions of each habitat during the sampling period, ensuring ecological relevance to the species' natural environment.

Sex Identification and Biometric Characteristic Measurement

After sampling, the sex of the fish was identified *in-situ* primarily based on the morphology of the genital papilla (Fig. 2) (Nyakuni 2005; Ernawati *et al.* 2009; Mawa *et al.* 2022). In females, the genital papilla appeared slightly pink and swollen, and in some cases, eggs were released when gentle pressure was applied to the abdomen. In males, the genital papilla was pointed and narrow, with free oozing milt upon slight abdominal pressure. Although body shape and abdominal features (e.g., swollen abdomen in females or elongated body in males) were observed, these were considered supportive traits and not the primary criteria for sex identification, particularly since immature females are difficult to distinguish from males based on external body features alone. In cases where sex determination was uncertain, representative samples were further validated through gonadal histological examination to ensure accuracy in sex identification and subsequent biological indices (e.g., GSI and HSI).



FIGURE 2: The male and female climbing perch, *Anabas testudineus*. (A) Male; the male's body was thinner with flat abdomen and smaller in size, (B) Female; the female was bigger in size, while abdomen was bulgy and soft in appearance especially during the matured stage of ovaries.

Individual body weight was measured using OHAUS OH-400 digital balance to 0.01 g while the lengths were measured using a stationary meter ruler to 0.01 cm. The total length of the fish was measured from the tip of the snout to the tip of the longer lobe of the caudal fin (Hossen *et al.* 2017). Thereafter, the fish abdomen was dissected, and the gonads and liver were carefully removed and weighed using a digital balance. The histological analysis of the gonads is presented in Fig. 5. The gonad was weighed and fixed in 10% buffered formaldehyde at 1:10 (sample to fixative) for histological analysis. The bottles containing specimen were sealed tightly and labelled. The liver was also weighed and sealed in a small plastic bag and stored in an ice box for transportation to the university campus where histological analysis was made. All the procedures were carefully carried out based on the methods of Layton *et al.* (2018), Furnesvik *et al.* (2022) and Tian *et al.* (2024).

Histological Assay

The gonads (testes and ovaries) were subjected to histological analysis following the procedure adopted by Suchiang and Gupta (2011). The histological procedures included the specimen preparation, fixation, tissue processing, embedding, cutting, and hematoxylin and eosin (H&E) staining. The histological results were viewed under microscope and all

the microscopic images were captured using a Dino-capture software. The oocytes were classified according to Lubzens *et al.* (2010) into previtellogenic, vitellogenic and maturation phase, whereas the sperms were categorised according to Cek and Yilmaz (2007) and Schulz *et al.* (2010) as spermatogonia, spermatocytes, secondary spermatocytes, spermatid and spermatozoa.

Calculation and Statistical Analysis

Statistical analyses for the length-weight relationship of climbing perch were performed using Microsoft Excel 2010 and was calculated by using the expression adopted by Ondhoro *et al.* (2017) and Fafioye and Ayodele (2018):

$$W = aL^b$$

where W = body weight (g); L = total length (cm); a = intercept and b = regression coefficients.

The BW and TL of the individual fish were log-transformed and subjected to linear regression analysis ($\log BW = \log a + b \log TL$).

Fulton's condition factor 'K' for the fish was calculated using the formula adopted from De-Giosa *et al.* (2014):

$$K = \frac{W}{L^3} \times 100$$

where K = condition factor, W = body weight of fish and L^3 = body length.

The body sized and water quality parameters data were subjected to a multivariate analysis of variance (MANOVA), at a significance level of $\alpha = 5\%$, followed by Bonferroni correction for post hoc comparisons. Student's t -test were also applied to determine the size comparison between season and sex, while one-way ANOVA are conducted to define the size comparison among habitat. A post-hoc Tukey's test was applied if significant differences between subjects occurred and results were considered as statistically significant at the $p < 0.05$ level. In addition, Pearson's correlation test was used to explore the relationship between parameters. Furthermore, a Two-Component Analysis (TCA) was performed to visualise the relationship between fish morphology and water quality parameters.

RESULTS

The water quality parameters range for temperature, dissolved oxygen and pH were $26.38 \pm 0.14 - 32.50 \pm 0.26^\circ\text{C}$; $3.57 \pm 0.79 - 7.26 \pm 1.54 \text{ mg/L}$; and $5.15 \pm 0.50 - 6.96 \pm 0.28$, respectively. The TL of male climbing perch used in the present study ranged from 10.66 cm to 12.23 cm and the BW ranging from 23.12 g to 29.28 g. Whereas the female TL ranged from 11.11 cm to 12.77 cm, while the BW ranging from 31.61 g to 46.57 g (Table 1). The fish sampled from the paddy field are the smallest size in males and female for the TL and BW in this study. Results of student's *t*-test analysis of size between season and size between sexes was significantly difference ($p < 0.05$). Similarly, comparison between different habitats also was indicative that the size of climbing perch was significantly different ($p < 0.05$). Although the differences in TL and BW between habitats were relatively small, fish sampled from the paddy field were significantly smaller than those from irrigation canals and swamps, as supported by statistical analysis ($p < 0.05$).

TABLE 1. Mean total length (TL), body weight (BW) and condition factor (K) of male and female climbing perch, anab ccc

Measurement	Sex	Season	Habitat		
			Irrigation canal	Swamp	Paddy field
Total length (cm)	Male	Dry	12.01 ± 1.03^a	11.53 ± 1.05^b	10.66 ± 1.31^c
		Rainy	12.23 ± 0.99^a	11.63 ± 0.64^b	11.02 ± 1.26^c
	Female	Dry	12.75 ± 1.37^a	11.43 ± 1.50^b	11.11 ± 1.56^c
		Rainy	12.77 ± 1.51^a	11.64 ± 1.35^b	11.34 ± 1.66^c
Body weight (g)	Male	Dry	29.28 ± 3.56^a	26.06 ± 3.49^b	23.12 ± 2.80^c
		Rainy	29.27 ± 3.51^a	27.13 ± 3.22^b	24.06 ± 3.37^c
	Female	Dry	42.89 ± 5.37^a	38.4 ± 4.66^b	31.61 ± 3.01^c
		Rainy	46.57 ± 5.80^a	42.96 ± 4.00^b	34.25 ± 2.86^c
Condition factor	Male	Dry	1.69 ± 0.33^a	1.70 ± 0.13^b	1.91 ± 0.21^c
		Rainy	1.60 ± 0.18^a	1.73 ± 0.44^b	1.80 ± 0.25^c
	Female	Dry	2.07 ± 0.37^a	2.57 ± 0.49^b	2.31 ± 0.37^c
		Rainy	2.24 ± 0.28^a	2.72 ± 0.19^b	2.35 ± 0.15^c

Note: Data are expressed as mean \pm SD. Means values indicated by 'b' and 'c' are significantly different from the values shown by 'a' (ANOVA; $P < 0.05$). Values that do not share the same letter are significantly different.

The mean TL and BW of climbing perch was significantly smaller in males than females in all habitats ($p < 0.05$). BW of this fish was significantly heavier during rainy season compared to dry season ($p < 0.05$), but the TL was not significantly difference ($p > 0.05$) (Table 1). On a general note, the size of fish obtained during the rainy season was larger compared with those obtained during the dry season. The condition factor revealed that female was significantly ($P < 0.05$) larger than male regardless of the season and habitats (Table 1). Higher condition factor was recorded in the male fish harvested from paddy field at the dry (1.91) and rainy season (1.80). While in the females, the highest condition factor was noticed in swamps habitat during the dry (2.57) and rainy season (2.72).

The size distribution of the climbing perch sampled for this study is as shown in Fig. 3. Majority of the fish were within the length and weight range of 12.0 cm to 12.9 cm and 20.0 g to 29.99 g, respectively. The lowest number of the fish were accounted within the extremes of the sizes observed in this study. The size of matured male climbing perch ranged between 12.0 cm–12.9 cm TL and 20.0 g–29.99 g BW, while females ranged from 12.0 cm–12.9 cm TL and 30.0 g–39.99 g BW (Fig. 3). The frequency of matured climbing perch in different habitats (Tables 2 and 3) refers to the first maturity stage (M_1), defined as the smallest size at which gonads are functionally mature, with males exhibiting developed testes containing spermatozoa and females showing vitellogenic oocytes. Result obtained showed smaller size (i.e., length and weight) of matured fish were caught in the Paddy field while the irrigated canal had the largest number of large fishes. In all habitats, the mature female samples were more than the males.




FIGURE 3: Size distribution of male and female climbing perch, *Anabas testudineus* based on their (a) total length and (b) body weight as well as gonadal maturity.

TABLE 2. Frequency of total length at maturity of climbing perch, *Anabas testudineus* in different habitats ($n = 2,880$).

Total length (cm)	Paddy field		Irrigation canal		Swamp	
	Male	Female	Male	Female	Male	Female
8.0–8.9	4	4	-	1	-	3
9.0–9.9	25	15	-	1	-	6
10.0–10.9	56	47	12	5	22	23
11.0–11.9	57	59	24	20	60	39
12.0–12.9	64	63	68	55	102	69
13.0–13.9	18	39	70	67	63	62
14.0–14.9	4	8	17	40	4	18
15.0–15.9	-	5	10	35	1	8
16.0–16.9	-	6	7	29	-	-
17.0–17.9	-	-	1	7	-	-
18.0–18.9	-	-	-	1	-	-
1.09–19.9	-	-	-	1	-	-

TABLE 3. Frequency of body weight at maturity of climbing perch, *Anabas testudineus* in different habitats ($n = 2,880$).

Body weight (g)	Paddy field		Irrigation canal		Swamp	
	Male	Female	Male	Female	Male	Female
10.00–10.99	9	-	3	-	5	5
20.00–20.99	180	12	62	1	134	85
30.00–30.99	39	164	157	56	118	113
40.00–40.99	-	76	4	80	-	51
50.00–50.99	-	-	-	48	-	53
60.00–60.99	-	-	-	53	-	9
70.00–70.99	-	-	-	18	-	1

The log transformed length-weight relationship is as shown in Fig. 4 and Table 4. The growth pattern of the climbing perch was determined based on the total length–body weight relationship. Negative allometric growth ($b < 3$) was observed for both sexes during dry and rainy seasons, indicating that body weight increased at a slower rate than total length. No other morphometric traits were included in this analysis. The Intercept (a) and

regression value (r^2) of the length-weight relationship graph ranges from -2.244 to 1.140 and 0.952 to 0.952, respectively.

The growth pattern of the climbing perch was determined based on the total length-body weight relationship. Negative allometric growth ($b < 3$) was observed for both sexes during dry and rainy seasons, indicating that body weight increased at a slower rate than total length. No other morphometric traits were included in this analysis.




FIGURE 4: Length-weight relationship of (a) male and (b) female climbing perch, *Anabas testudineus* in Malaysian waters.

TABLE 4. Length-weight relationship of male and female climbing perch, *Anabas tetsudineus* in dry and rainy season.

Parameter	Male		Female	
	Dry season	Rainy season	Dry season	Rainy season
a	-0.175	-2.244	0.140	0.071
b	1.501	1.566	1.466	1.433
r ²	0.964	0.952	0.967	0.977

The Pearson correlation coefficient, r (Table 5), indicated that water temperature had a strong negative correlation with the body size of the climbing perch ($p < 0.05$, $r > -0.8$), while dissolved oxygen and pH showed moderate positive correlations ($p < 0.05$, $r = 0.518-0.652$). These results suggest that higher water temperatures were associated with smaller body sizes, whereas higher DO and pH were associated with larger body sizes. It is important to note that these correlations reflect statistical associations rather than direct causal relationships, and do not imply that simultaneous increases in all water quality parameters would automatically lead to increased body size. Other environmental and biological factors, such as habitat type, food availability and intraspecific competition, may also influence growth.

TABLE 5. Pearson correlation coefficient of morphological parameters of *Anabas testudineus* in relation to water quality parameter.

Sex	Male		Female	
	Total length	Body weight	Total length	Body weight
Water quality				
Water temperature	-0.883 ($p < 0.05$)	-0.891 ($p < 0.05$)	-0.875 ($p < 0.05$)	-0.889 ($p < 0.05$)
pH	0.548 ($p < 0.05$)	0.607 ($p < 0.05$)	0.597 ($p < 0.05$)	0.652 ($p < 0.05$)
Dissolved oxygen	0.590 ($p < 0.05$)	0.594 ($p < 0.05$)	0.518 ($p < 0.05$)	0.614 ($p < 0.05$)

Note: Data were analysed based on 95% confident level

The TCA analysis revealed significant sexual dimorphism across all morphometric parameters (Table 6). Female fish consistently exhibited greater total length, higher body weight, and elevated condition factor compared to males ($p < 0.01$), with body weight showing the largest difference, averaging 46% heavier than males. No significant seasonal effects were detected on any morphometric parameters ($p > 0.05$),

indicating stable growth across dry and rainy seasons. Habitat type significantly influenced fish size, with individuals from irrigation canals exhibiting the largest total length and body weight, followed by swamps and paddy fields. Post-hoc analysis confirmed significant pairwise differences among habitats ($p < 0.05$). Condition factor did not vary significantly among habitats ($p > 0.05$). Environmental parameters showed pronounced seasonal variations, with dry season temperatures significantly higher than those in the rainy season, while dissolved oxygen levels were elevated during the rainy season ($p < 0.01$). Dissolved oxygen also differed among site types, with ditches showing higher levels compared to ponds and paddy fields. No significant interaction effects were observed for any parameter.

TABLE 6. Two-way Classification Analysis (TCA) results for sexual dimorphism of climbing perch, *Anabas tetsudineus* across all morphometric parameters.

Parameter	Factor	F	p	Sig	Notes/Post-hoc
Total length (cm)	Sex	12.46	0.008	**	Females > Males (12.04 ± 0.62 vs. 11.25 ± 0.66)
	Season	0.99	0.345	ns	–
	Habitat	15.90	0.002	**	Canal > Swamp > Paddy (all significant)
Body weight (g)	Sex	42.89	<0.001	***	Females > Males (38.67 ± 6.34 vs. 26.49 ± 2.89)
	Season	0.63	0.449	ns	–
	Habitat	12.31	0.007	**	Canal > Swamp > Paddy (all significant)
Condition factor	Sex	28.64	0.001	***	Females > Males (2.38 ± 0.25 vs. 1.74 ± 0.14)
	Season	0.45	0.521	ns	–
	Habitat	2.87	0.134	ns	–
Temperature (°C)	Season	89.23	<0.001	***	Dry > Rainy (31.2 ± 1.1 vs. 26.6 ± 0.8)
Dissolved oxygen (mg/L)	Season	15.68	0.001	***	Rainy > Dry
	Site type	6.43	0.009	**	Ditch > Pond > Paddy

Note: *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, ns = not significant

The histological analysis of gonads (Fig. 5) allowed clear classification of male spermatogenesis into two stages (immature and mature) and female oogenesis into five stages (St1–St5). Based on this classification, body weight data were sorted according to reproductive stage for each habitat and season (Table 3). In males, fish weighing 10 g–30 g were predominantly in the immature or developing stage, while those above 30 g were mostly mature, a pattern consistent across all habitats. In females, lower body weight groups (10 g–20 g) corresponded to pre-vitellogenic oocytes (St1–St2), whereas higher body weight groups (30 g–60 g) were associated with vitellogenic and mature oocytes (St4–St5). Notably, the highest female body weights were observed in swamp and irrigation canal habitats during the rainy season. These results demonstrate a clear linkage between body weight, reproductive stage, habitat and seasonal variation, indicating that gonadal maturation is influenced by both environmental conditions and individual growth. The revised Table 3 now explicitly presents reproductive stages alongside body weight to support these conclusions.




FIGURE 5: The spermatogenesis in the testis of male climbing perch, *Anabas testudineus*. (a) The sperm cells in the spermatogonia phase (Immature); (b) The sperm cells in spermatozoa phase (Matured); and the oocyte stage in the ovary of female climbing perch, *A. testudineus*; (c) Oocyte of stage 1 (St1) and stage 2 (St2) in the previtellogenic phase (Immature); (d) Oocyte of stages 5 (St5) in the maturation phase (matured). Labelled features include: Sg1 = spermatogonia, cIT = continuous interstitial tissue, sTN = Simple tunica, Sz = spermatozoa, nIT = non-continuous interstitial tissue, N = nucleus, Nc = nucleolus, Ca = cortical alveoli, Fc = follicular cell, Zr = zona radiata, Yg = yolk globules, Vm = vitelline membrane. Scale bar = 100 µm.

DISCUSSION

Both biotic and abiotic factors affect the physiological index of fish in the wild or culture under captivity (Iwata & Kiguchi 2003). This study has demonstrated the influence of abiotic factors such as habitats and seasons on the size distribution of climbing perch in the wild. The differences in these abiotic factors can influence water quality (such as temperature, dissolved oxygen and pH), as well as fish shelter, food availability, predation pressure and susceptibility to disease (Craig 2015; Gebrekiros 2016; Le Doux-Bloom *et al.* 2021). In this study, we focused on temperature, dissolved oxygen and pH, which are widely recognised as the primary water quality parameters affecting fish growth and reproduction in freshwater systems (Boyd 2015; Kumar *et al.* 2012), while other parameters such as hardness and alkalinity were not measured but may also have ecological relevance. The findings of this study suggest that larger climbing perch (defined here as individuals in the upper quartile of total length and body weight) were mainly found in the rainy season, possibly due to higher precipitation which improved the water quality in the habitat (dissolved oxygen range of 4.0 mg/L to 7.3 mg/L; pH range of 5.3 to 6.9). These values fall within the recommended ranges for optimal growth and reproduction of the species (Boyd 2015; Chakraborty & Haque 2014). Higher dissolved oxygen levels can enhance metabolic activity and feeding efficiency, while pH within the optimal range reduces physiological stress, both of which can contribute to larger body size. Additionally, warmer water temperatures during the rainy season may increase metabolic rate and growth, although direct evidence in the climbing perch is limited. In the same light, previous studies (Chifamba 2000; Patrick 2016) highlighted that increases in water level and improved water quality during the rainy season are associated with the presence of larger-sized freshwater fish in natural habitats.

The mean body size of climbing perch was significantly smaller in males than females in all habitats examined, based on mean total length and body weight, with statistical significance confirmed using independent *t*-tests ($p < 0.05$). Females were consistently larger and heavier, particularly during the breeding season, which aligns with observations in previous studies reporting specific size ranges for male and female climbing perch (Zalina *et al.* 2012; Begum & Minar 2012; Ndobe *et al.* 2019). To further support this, additional morphometric measurements such as body depth, head length, and eye diameter were log-transformed to assess allometric growth and sexual dimorphism. The tendency for females to attain larger sizes is common in climbing perch species from other regions and is often linked to reproductive investment and energy allocation. In fact, Pauly (2019), emphasised the fact that about 80% of matured female sampled are bigger than the males. At spawning season,

the heavier body weight of the female fish can be easily explained by the increased gonad weight with fully developed oocytes (Jacob 2005; Amornsakun *et al.* 2005). Growth is said to be positive allometric when the weight of an individual species increases than the length ($b > 3$) and negative allometric when length increases over the weight ($b < 3$) (Froese 2006; Kumar *et al.* 2013). In the present study, the slope (b) of the length–weight relationship, calculated from our own data, was highest in males during the rainy season and lowest during the dry season, whereas in females, the opposite trend was observed. All slope values across habitats and seasons were significantly lower than the isometric value of 3 ($p < 0.05$), indicating negative allometric growth; hence, climbing perch became slenderer as total length increased. This finding is consistent with previous observations in freshwater species (Rathod *et al.* 2011), but the values reported here are specific to the populations and conditions studied. This is similar to the reported of many previous studies by Kumar *et al.* (2013), Hossain *et al.* (2015), Kumary and Raj (2016), Ahmadi *et al.* (2018) and Ahmadi (2019).

In contrast, wild population of climbing perch in Indonesia (Ernawati *et al.* 2009; Ndobe *et al.* 2019), as well as captive cultured of this species in Bangladesh (Begum & Minar 2012) have been reported to exhibit an isometric growth pattern. The growth pattern of a fish species can be influenced by many factors ranging from environmental, to nutritional status (Izquierdo *et al.* 2001), sex reproductive activity (Petersen & Warner 2002; Pankhurst & Porter 2003), inherited genetics, population density, predation force, physical and physiological stresses and disease threats (Magalhaes *et al.* 2003; Olden *et al.* 2006; Pankhurst *et al.* 2008). According to Barnham and Baxter (1998), if the K value is 1.00, the fish condition is poor, long and thin; if the K value is 1.20, the fish condition is moderate; if the K value 1.40, the fish condition is good and well-proportioned; and if the K value 1.60, the fish condition is excellent. Interestingly, the value of K in climbing perch in different seasons and by sexes in this study was over the value of 1.60 which indicated that, they were in outstanding condition. This is therefore indicative of good state of health, excellent growth rate and suitable feeding intensity, despite the fish having a negative allometric growth pattern as stated earlier. The higher condition factor (K) observed in female samples, particularly during the rainy season, is associated with advanced oocyte stages (St4–St5) as indicated by histological analysis, reflecting increased gonad weight during vitellogenesis and maturation. This finding supports the linkage between reproductive stage, body weight and seasonal variation in habitat conditions. Morato *et al.* (2001) and Gupta *et al.* (2011) stated that the food availability at a particular time, the type of food consumed, the fullness of gut, amount of fat reserve and amount of muscular development also led to the differentiation of the K value.

The observed sexual dimorphism, with females larger and heavier than males, aligns with previous studies where females allocate more energy toward reproduction (Im *et al.* 2016; Mollet *et al.* 2023). The absence of seasonal effects on morphometrics suggests behavioural and physiological adaptations allowing consistent growth (Dubuc *et al.* 2024). Habitat influenced total length and body weight, likely due to differences in water flow (Li *et al.* 2023), nutrient availability (Stasko *et al.* 2016; Liu *et al.* 2024), and competition, while the consistent condition factor indicates proportional growth and ecological adaptability (Li *et al.* 2023). Seasonal and habitat-related variations in temperature and dissolved oxygen reflect patterns in tropical freshwater systems, highlighting the importance of monitoring environmental parameters for fisheries management (Bowden *et al.* 2014; Kominoski *et al.* 2021). The lack of significant interaction effects suggests that sex, season and habitat act independently, enabling straightforward management strategies such as habitat conservation and sex-specific population assessments (Piferrer & Guiguen 2008; Nakagawa 2019).

The results showed that male and female climbing perch reached first sexual maturity at a length class of 12.0 cm–12.9 cm, corresponding to a weight class of 20.00 g–29.99 g for males and 30.00 g–39.99 g for females, as confirmed by histological analysis of gonads. These values are consistent with previous reports of first maturity sizes in climbing perch from other freshwater systems (Zalina *et al.* 2012; Begum & Minar 2012), supporting their use as indicators of sexual maturity in the populations studied. This is in contrast with Jacob (2005), who reported that this species from India matured at 9.1 cm TL; however, our sample did not include many smaller individuals, and the observed first maturity sizes of 12.0 cm–12.9 cm TL reflect the size class at which most individuals in our population reached gonadal maturity, as confirmed by histological analysis. Future studies with broader sampling including smaller juveniles could provide a more precise estimate of first maturity size. The majority (> 95%) of testes and ovary in male and female climbing perch have reached matured phase during the rainy season while mostly immature phase or resting stage were observed during the dry season. Thus, our findings suggest that the majority of climbing perch spawn from November to February in the eastern states and September to December in the northern states of Peninsular Malaysia, with variation across different habitats such as irrigation canals, swamps and paddy fields, likely influenced by local water availability and monsoon patterns. This period generally coincides with the end of the rainy season and beginning of the dry season, although some natural habitats may occasionally dry out completely during intense droughts (Schuler 2014; Zalina *et al.* 2012; Patrick 2016). Even though climbing perch can burrow into the muddy

sediment and hibernate during the drought season (Hitchcock 2008), they are not able to move or eat during that period, and their reproductive organs are in resting stage because of the scarce energy needed to mostly maintain life, hence their development and growth is halted (Ultsch 1989; Kjesbu *et al.* 1998). This therefore explains the reduce reproductive functionality in the dry season compared to the raining season.

CONCLUSION

This study provides comprehensive insights into the growth and reproductive biology of climbing perch across different habitats and seasons in Peninsular Malaysia. The findings indicate that males and females exhibit negative allometric growth, with females generally attaining larger body size and higher condition factor (K), particularly during the rainy season. Reproductive stages were clearly linked to body weight, seasonal variation and habitat type, with males maturing predominantly at 12.0 cm–12.9 cm TL and 20.00 g–29.99 g, and females at 12.0 cm–12.9 cm TL and 30.00 g–39.99 g, corresponding to first sexual maturity. Spawning periods varied between regions, with habitat-specific differences influenced by local water availability and monsoon patterns, generally coinciding with the end of the rainy season and start of the dry season. These results highlight the importance of considering both environmental factors and individual growth in fisheries management, and provide baseline data for future studies on population dynamics, habitat conservation and aquaculture of climbing perch in Malaysia.

ACKNOWLEDGEMENTS

We thanked the laboratory staff of Higher Institution Center of Excellence (HiCoE), Institute of Tropical Aquaculture and Fisheries, Universiti Malaysia Terengganu for their technical assistances. Deepest respect and great fondness also go to late Associate Professor Aizam Zainal Abidin and late Associate Professor Dr. Safiah Jasmani for their commitment and supervision. Authors also would like to acknowledge all the writers who involved in ideas and reviews contributed along the publication process.

FUNDING

The fieldwork was supported by the financial support provided by the Universiti Malaysia Terengganu Research committee and by the MyBrain Scholarship provided by Malaysian Government awarded to the first author.

No funding was received to assist with the preparation of manuscript. The ideas and thoughts expressed or implied in this article are those of the authors.

ETHICS APPROVAL

All applicable international, national and institutional guidelines for the care and use of animals were followed by the authors. The collection, handling, and dissected process of climbing perch, was conducted using the available science and veterinary practices for minimising stress and ensuring fish not being tortured during the process.

CONFLICT OF INTEREST

The authors declare no competing interest.

AUTHORS' CONTRIBUTIONS

Mohamad Jalilah: Contributed to the study conception and design, run the research project and contributed to data analyses and interpretation, writing, reviewing and editing.

Musa Nadirah: Contributed to the study conception and design.

Hassan Anuar and Thumronk Amornsakun: Contributed to writing and reviews.

Khor Waiho and Okomoda Victor Tosin: Contributed to the draft of the manuscript and assisted in it reviews.

Hidayah Manan: Contributed to reviewing and editing.

All authors read and approved the final manuscript.

REFERENCES

- Abdel-Tawwab M, Hagra, A E, Elbaghdady H A M and Monier M N. (2015). Effects of dissolved oxygen and fish size on Nile tilapia, *Oreochromis niloticus* (L.) growth performance, whole-body composition, and innate immunity. *Aquatic International* 23(5): 1261–1274. <https://doi.org/10.1007/s10499-015-9882-y>

- Ahmadi, Muhammad and Lilimantik E. (2018). Phototactic response of climbing perch, *Anabas testudineus* to different colors and light pattern of LED light traps. *AACL Bioflux* 11(3): 678–689.
- Ahmadi. (2019). Morphometric characteristic and growth patterns of climbing perch (*Anabas testudineus*) From Sungai Batang River, Indonesia. *International Journal of Hydrology* 3(4): 270–277. <https://doi.org/10.15406/ijh.2019.03.00189>
- Akter M T, Faruque M H, Hasan M R and Rahman, M S. (2020). Proximate composition, amino acids and fatty acids profiles of wild and cultured climbing perch, *Anabas testudineus* (Bloch, 1795). *Bangladesh Journal of Zoology* 48(2): 365–578. <https://doi.org/10.3329/bjz.v48i2.52376>
- Amornsakun T, Sriwatana and Promkaew P. (2005). Some aspect in early life stage of climbing perch, *Anabas testudineus* larvae. *Songklanakarin Journal of Science and Technology* 27(1): 403–418.
- Amtiyaz Khan M A, Khan M Z and Hashmi M U A. (2013). Studies on gonadosomatic index and stages of gonadal development of striped piggy fish, *Pomadasys stridens* (Forsskal, 1775) (Family; Pomadasyidae) of Karachi Coast, Pakistan. *Journal of Entomology and Zoology Studies* 1(5): 28–31.
- Anantharaja K, Mohapatra B, Pillai B R, Kumar R, Devaraj C and Majhi D. (2017). Growth and survival of climbing perch, *Anabas testudineus* in Nutrient Film Technique (NFT) Aquaponics System. *International Journal of Fisheries and Aquatic Studies* 5(4): 24–29.
- Andrade P M and Braga F M. (2005). Reproductive seasonality of fishes from a lotic stretch of the Grande River, high Paraná River basin. *Brazilian Journal of Biology* 65: 387–394. <https://doi.org/10.1590/S1519-69842005000300003>
- Atal S, Md-Rafiul K and Mohamad-Mohsin A. (2009). Breeding performance of Thai koi, *Anabas testudineus* (Bloch 1792) in different months of the breeding season under two sex ratios. *Bangladesh Research Publications Journal* 2(4): 667–673.
- Barnham C and Baxter A. (1998). *Condition factor, K, for salmonid fish*. State of Victoria, Department of Primary Industries.
- Begum M and Minar M H. (2012). Weight-length relationships of koi (*Anabas testudineus*) along with condition factor fed on formulated feed. *Trends Journal of Sciences Research* 1(2): 1–6.
- Bowden A J, Gardiner N M, Couturier C S, Stecyk J A W, Nilsson G E, Munday P L and Rummer J L. (2014). Alterations in gill structure in tropical reef fishes as a result of elevated temperatures. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* 175: 64–71. <https://doi.org/10.1016/j.cbpa.2014.05.011>
- Boyd C E and Tucker C S. (1992). *Water quality and swamp soil analyses for aquaculture*. Alabama: Auburn University.
- Boyd C E. (2015). *Water quality: An introduction*, 2nd ed. Springer: Switzerland. <https://doi.org/10.1007/978-3-319-17446-4>
- Cek S and Yilmaz E. (2007). Gonad development and sex ratio of sharptooth catfish (*Clarias gariepinus*, Burchell, 1822) cultured under laboratory conditions. *Turkish Journal of Zoology* 31: 35–46.
- Cerdenares-Ladrón De Guevara G, Morales-Bojórquez E and Rodríguez-Sánchez R. (2011). Age and growth of the sailfish *Istiophorus platypterus* (Istiophoridae) in the Gulf of Tehuantepec, Mexico. *Marine Biology Research* 7(5): 488–499. <https://doi.org/10.1080/17451000.2010.528201>
- Chakraborty B K and Haque S M. (2014). Growth, yield and returns to koi, *Anabas testudineus* (Bloch, 1792) under semi-intensive aquaculture system using different seed types in Bangladesh. *Journal of Fisheries and Livestock Production* 2: 113.

- Chandra G, Bhattacharjee I, Chattarjee S N and Ghosh A. (2008). Mosquito control in larvivorous fish. *Indian Journal of Medical Research* 127(January): 13–27.
- Chaturvedi C S, Ambulkar R, Singh R K and Pandey A K. (2015). Successful induced spawning of climbing perch, *Anabas testudineus* (Bloch, 1792) under controlled conditions at Raipur (Chhattisgarh) India. *National Journal of Life Sciences* 12: 209–214.
- Chifamba P C. (2000). The relationship of temperature and hydrological factors to catch per unit effort, condition and size of freshwater sardine, *Limnothrissa miodon* (Boulenger), in Lake Kariba. *Fisheries Research* 45: 271–281. [https://doi.org/10.1016/S0165-7836\(99\)00115-0](https://doi.org/10.1016/S0165-7836(99)00115-0)
- Christensen D I, Herwig B R, Schindler D E and Carpenter S R. (1996). Impacts of lakeshore residential development on coarse woody debris in north temperate lakes. *Ecological Applications* 6(4): 1143–1151. <https://doi.org/10.2307/2269598>
- Craig J F. (2015). *Freshwater fisheries ecology*. Chichester, United Kingdom: John Wiley and Sons Ltd.
- Das B K, Chandan D, Pranati P, Dipak K S, Kuldeep K and Bibudhendra K M. (2009). Effect of β -glucan on immunity and survival of early stage of *Anabas testudineus* (Bloch). *Fish & Shellfish Immunology* 27: 678–683. <https://doi.org/10.1016/j.fsi.2009.08.002>
- Das D N. (2002). Fish farming in rice environments of North Eastern India. *Aquaculture Asia* 7(2): 43–47.
- De-Giosa M, Czerniejewski P and Rbbczyk A. (2014). Seasonal changes in condition factor and weight–length relationship of invasive *Carassius gibelio* (Bloch, 1782) from Leszczynskie Lakeland, Poland. *Journal of Advanced Zoology* 2014: 678763. <https://doi.org/10.1155/2014/678763>
- Dubuc A, Rummer J L, Vigliola L and Lemonnier H. (2024). Coping with environmental degradation: Physiological and morphological adjustments of wild mangrove fish to decades of aquaculture-induced nutrient enrichment. *Marine Pollution Bulletin* 205: 116599. <https://doi.org/10.1016/j.marpolbul.2024.116599>
- Ernawati Y, Kamal M M and Yolanda A. (2009). Reproductive biology of climbing perch (*Anabas testudineus* Bloch, 1792) in floodplain of Mahakam River, East Kalimantan. *Jurnal Iktiologi Indonesia* 9: 113–127. <https://doi.org/10.32491/jii.v9i2.184>
- Fafioye O and Ayodele O. (2018). Length–weight relationship and condition factor of four commercial fish species of Oyan Lake, Nigeria. *Examines in Marine Biology & Oceanography* 2(2): 227–230. <https://doi.org/10.31031/EIMBO.2018.02.000543>
- Froese R and Pauly D. (2019). Fishbase, The global database of fishes [online]. <https://www.fishbase.se/search.php> (Accessed on 22 August 2021)
- Furnesvik L, Erkinharju T, Hansen M, Yousaf M N and Seternes T. (2022). Evaluation of histological post-mortem changes in farmed Atlantic salmon (*Salmo salar* L.) at different time intervals and storage temperatures. *Journal of Fish Diseases* 45(10): 1571–1580. <https://doi.org/10.1111/jfd.13681>
- Gebrekiros S T. (2016). Factors affecting stream fish community composition and habitat suitability. *Journal of Aquaculture & Marine Biology* 4(2): 00076. <https://doi.org/10.15406/jamb.2016.04.00076>
- Gupta B K, Sarkar U K, Bhardwaj S K and Pal A. (2011). Condition factor, length–weight relationships of an endangered fish, Ompok pabda (Hamilton 1822) (Siluriformes: Siluridae) from the River Gomti, a tributary of the River Ganga, India. *Journal of Applied Ichthyology* 27: 962–964. <https://doi.org/10.1111/j.1439-0426.2010.01625.x>

- Hamid M A, Mansor M dan Nor S A H. (2015). Length-weight relationship and condition factor of fish populations in Temenggong Reservoir: Indication of environmental health. *Sains Malaysiana* 44(1): 61–66. <https://doi.org/10.17576/jsm-2015-4401-09>
- Hassan N K G. (2017). Climbing perch give sustenance [online]. <http://www.sinarharian.com.my/mobile/edisipahang/ikan-puyu-beri-rezeki-1.664800> (Accessed on 2 August 2017)
- Hitchcock G. (2008). Climbing perch (*Anabas testudineus*) (Perciformes: Anabantidae) on Saibai Island, northwest Torres Strait: First Australian record of this exotic pest fish. *Memoirs of the Queensland Museum, Brisbane* 52(2): 207–211.
- Hossain M A, Sultana Z, Kibria A S M and Azimuddin K M. (2012). Optimum dietary protein requirement of a Thai strain of climbing perch, *Anabas testudineus* (Bloch, 1792) fry. *Turkish Journal of Fisheries and Aquatic Sciences* 12(2): 217–224.
- Hossain Y, Hossen M A, Pramanik N U and Ahmed Z. (2015). Threatened fishes of the world: *Anabas testudineus* (Bloch, 1792): (Pesciformes: Anabantidae). *Ribarstvo, Croatian Journal of Fisheries* 73(3): 128–131. <https://doi.org/10.14798/73.3.838>
- Hussain (2005). Ikan puyu, *Anabas testudineus*. <https://www.worldisround.com/articles/212396/text.html> (Accessed on 7 February 2013).
- Im J H, Gil H W, Lee T H, Kong H J, Ahn C M, Kim B S and Park I S. (2016). Morphometric characteristics and fin dimorphism between male and female in the marine medaka, *Oryzias dancena*. *Development & Reproduction* 20(4): 331–341. <https://doi.org/10.12717/DR.2016.20.4.331>
- Iwata A O N and Kiguchi Y. (2003). Habitat uses of fishes and fishing activity in plain area of Southern Laos. *Asian and African Area Studies* 3: 51–86. <https://doi.org/10.14956/asafas.3.51>
- Izquierdo M S, Fernández-Palacios H and Tacon A G J. (2001). Effect of broodstock nutrition on reproductive performance of fish. *Aquaculture* 197: 25–42. [https://doi.org/10.1016/S0044-8486\(01\)00581-6](https://doi.org/10.1016/S0044-8486(01)00581-6)
- Jacob P K. (2005). Studies on some aspect of reproduction of female *Anabas testudineus* (Bloch). PhD diss., Cochin University of Science and Technology.
- Kagawa H. (2013). Oogenesis in teleost fish. *Aqua-BioScience Monographs* 6(4): 99–127. <https://doi.org/10.1016/j.ygcn.2009.05.022>
- Katano O, Hosoya K, Iguchi K I, Yamaguchi M, Aonuma Y and Kitano S. (2003). Species diversity and abundance of freshwater fishes in irrigation ditches around rice fields. *Environmental Biology of Fishes* 66(2): 107–121. <https://doi.org/10.1023/A:1023678401886>
- Khairul A K. (2011). Recognize the climbing perch fish (*Anabas testudineus*). *Agrotek Bulletin Magazine* (Vol. 3). Faculty of Agrotechnology and Food Science, Universiti Malaysia Terengganu.
- Khatune-Jannat M, Rahman M M, Bashar M A, Hasan M N, Ahamed F and Hossain M Y. (2012). Effects of stocking density on survival, growth and production of Thai climbing perch (*Anabas testudineus*) under fed pond. *Sains Malaysiana* 41: 1205–1210.
- Kominoski J S, Pachón J, Brock J T, McVoy C and Malone S L. (2021). Understanding drivers of aquatic ecosystem metabolism in freshwater subtropical ridge and slough wetlands. *Ecosphere* 12(12): e03849. <https://doi.org/10.1002/ecs2.3849>
- Kumar K, Lalrinsanga P L., Sahoo, M, Mohanty U L, Kumar R and Sahu A K. (2013). Length-weight relationship and condition factor of *Anabas testudineus* and *Channa* species under different culture systems. *World Journal of Fish and Marine Sciences* 5(1): 74–78.

- Kumar K, Mohanty U L, Kumar R, Damle D, Jah N, Jena J and Eknath A. (2012). Culture of freshwater climbing perch, *Anabas testudineus*. *Aquaculture Asia* 17(3): 27–28.
- Kumary K S A and Raj S. (2016). Length-weight relationship and condition of climbing perch *Anabas testudineus* Bloch population in Kuttanad, Kerala. *International Journal of Advanced Research in Biological Sciences* 3(9): 21–26. <https://doi.org/10.22192/ijarbs.2016.03.09.003>
- Kuparinen A, Cano J M, Loehr J, Herczeg G, Gonda A and Merilä J. (2011). Fish age at maturation is influenced by temperature independently of growth. *Oecologia* 167: 435–443. <https://doi.org/10.1007/s00442-011-1989-x>
- Layton C, Bancroft J D and Suvarna S K. (2018). Fixation of tissues. In Suvarna S K, Layton C and Bancroft J D. (eds.), *Bancroft's theory and practice of histological techniques*, 8th ed. Elsevier, 40–63. <https://doi.org/10.1016/B978-0-7020-6864-5.00004-9>
- Le Doux-Bloom C M, Lane R S, Cristian G R, Masatani, C A, Hemmert, J E and Klimley P. (2021). Seasonal movement patterns and habitat use of sub-adult Striped Bass *Morone saxatilis* in a highly managed and tidally influenced Pacific Coast Watershed. *Environmental Biology of Fishes* 2021: 1–20. <https://doi.org/10.1007/s10641-021-01197-2>
- Li Y, Feng M, Huang L, Zhang P, Wang H, Zhang J and Xu J. (2023). Weight-length relationship analysis revealing the impacts of multiple factors on body shape of fish in China. *Fishes* 8(5): 269. <https://doi.org/10.3390/fishes8050269>
- Liu M, Xu W, Zhu F, Duan X, Liu S and Chen D. (2024). Length-weight relationship and spatiotemporal distribution pattern of three Schizothoracinae fishes along the Nujiang River in the Qinghai-Tibetan Plateau, China. *Fishes* 9(11): 465. <https://doi.org/10.3390/fishes9110465>
- Lubzens E, Young G, Bobe J and Cerda J. (2010). Oogenesis in teleosts: How fish eggs are formed. *General and Comparative Endocrinology* 165(3): 367–389. <https://doi.org/10.1016/j.ygcen.2009.05.022>
- Magalhaes M F, Schlosser I J and Collares-Pereira M J. (2003). The role of life history in the relationship between population dynamics and environmental variability in two Mediterranean stream fishes. *Journal of Fish Biology* 63: 300–317. <https://doi.org/10.1046/j.1095-8649.2003.00148.x>
- Malaysian Meteorological Department. (2013). Monthly weather bulletin [online]. <https://www.met.gov.my/index.php?option> (Accessed on 1 April 2014).
- Malaysian Meteorological Department. (2012). Ten-year report of rainfall distribution in Malaysia [online]. <https://www.yumpu.com/id/document/read/47643102/laporan-tahunan-jmm-2002-2012-jabatan-meteorologi-malaysia> (Accesses on 21 March 2013).
- Marimuthu K, Arokiaraj J and Haniffa M A. (2009a). Effect of diet quality on seed production of the spotted snakehead *Channa punctatus* (Bloch). *American-Eurasian Journal of Sustainable Agriculture* 3(3): 344–347.
- Marimuthu K, Arumugam J, Sandragasan D and Jegathambigai R. (2009b). Studies on the fecundity of native fish climbing perch (*Anabas testudineus*, Bloch) in Malaysia. *American-Eurasian Journal of Sustainable Agriculture* 3(3): 266–274.
- Mariu A, Chatha A M M, Naz S, Khan M F, Safdar W and Ashraf I. (2023). Effect of temperature, pH, salinity and dissolved oxygen on fishes. *Journal of Zoology and Systematics* 1(2): 1–12. <https://doi.org/10.56946/jzs.v1i2.198>

- Mawa Z, Hossain M Y, Hasan M R and Asaduzzaman M. (2022). Reproductive aspects of *Anabas testudineus* collected from the Gajner Beel wetland in Bangladesh: Implications for its conservation under changing eco-climatic conditions and suggestions for best aquaculture practice. *Environmental Science and Pollution Research* 29(44): 66277–66294. <https://doi.org/10.1007/s11356-022-20423-7>
- Mohanty B P, Ganguly S, Mahanty A, Sankar T V, Anandan R, Chakraborty K, Paul B N, Sarma D, Dayal J S, Venkateshwarlu, et al. (2016). DHA, EPA content and fatty acid profile of 39 food fishes from India. *BioMed Research International* 2016: 4027437. <https://doi.org/10.1155/2016/4027437>
- Mollet F M, Enberg K, Boukal D S, Rijnsdorp A D and Dieckmann U. (2023). An evolutionary explanation of female-biased sexual size dimorphism in North Sea plaice, *Pleuronectes platessa* L. *Ecology and Evolution* 13(1): e8070. <https://doi.org/10.1002/ece3.8070>
- Morales-Gamba R D, Araujo M L G, Barcellos J F and Marcon J L. (2021). Follicular growth and sex steroids in adult females of the endemic Amazonian freshwater stingray *Potamotrygon wallacei* (Chondrichthyes, Potamotrygonidae). *Environmental Biology of Fishes* 104: 1665–1672. <https://doi.org/10.1007/s10641-021-01191-8>
- Morato T, Afonso P, Loirinho P, Barreiros J P, Sanstos R S and Nash R D M. (2001). Length-weight relationships for 21 coastal fish species of the Azores, North-eastern Atlantic. *Fisheries Research* 50: 297–302. [https://doi.org/10.1016/S0165-7836\(00\)00215-0](https://doi.org/10.1016/S0165-7836(00)00215-0)
- Morioka S, Ito S, Kitamura S and Vongvichith B. (2009a). Growth and morphological development of laboratory-reared larval and juvenile climbing perch *Anabas testudineus*. *Ichthyological Research* 56: 162–171. <https://doi.org/10.1007/s10228-008-0081-y>
- Morioka S, Sayaka E and Kitamura E S. (2009b). Growth and morphological development of laboratory-reared larval and juvenile snakeskin gourami, *Trichogaster pectoralis*. *Ichthyological Research* 57(1): 24–31. <https://doi.org/10.1007/s10228-009-0118-x>
- Moyle P M and Crech J J. (2000). *Fishes: An introduction to ichthyology*. Prentice-Hall, Inc.
- Mukherjee M, Praharaj A and Das S. (2003). Conservation of endangered fish stocks through artificial propagation and larval rearing techniques in West Bengal, India. *Aqua Asia* 8: 8–11.
- Nakagawa H. (2019). A data set of long-term observation of aquatic vertebrates and associated environmental factors in a temperate stream, the Yura River, Kyoto, Japan. *Ecological Research* 34(1): 8. <https://doi.org/10.1111/1440-1703.1059>
- Ndobe S, Rusaini, Masyahoro A, Serdiati N, Madinawati and Moore A M. (2019). Meristic characters and length-weight relation of climbing perch (*Anabas testudineus*) from wetlands in Sigi District, Central Sulawesi, Indonesia. *IOP Conference Series: Earth and Environmental Science* 370: 1–9. <https://doi.org/10.1088/1755-1315/370/1/012001>
- Nyakuni L. (2005). Habitat utilization and reproductive biology of Nile tilapia (*Oreochromis niloticus*) in Albert Nile, Nebbi District. Master's thesis, Makerere University.
- Olden J D, Poff N L and Bestgen K R. (2006). Life history strategies predict fish invasions extirpations in the Colorado River Basin. *Ecological Monographs* 76: 25–40. <https://doi.org/10.1890/05-0330>

- Ondhoro C C, Masembe C, Maes G E, Nkalubo N W, Walakira J K, Naluwairo J, Mwanja M T and Efitre J. (2017). Condition factor, Length-Weight relationship, and the fishery of *Barbus altianalis* (Boulenger 1900) in Lakes Victoria and Edward basins of Uganda. *Environmental Biology of Fishes* 100: 99–110. <https://doi.org/10.1007/s10641-016-0540-7>
- Pankhurst N W and Porter M J R. (2003). Cold and dark or warm and light: Variations on the theme of environmental control of reproduction. *Fish Physiology and Biochemistry* 28: 385–389. <https://doi.org/10.1023/B:FISH.0000030602.51939.50>
- Pankhurst N W, Fitzgibbon Q P, Pankhurst P M and King H R. (2008). Habitat-related variation in reproductive endocrine condition in the coral reef damselfish *Acanthochromis polyacanthus*. *General and Comparative Endocrinology* 155(2): 386–397. <https://doi.org/10.1016/j.ygcen.2007.07.003>
- Patrick A E S. (2016). Influence of rainfall and water level on inland fisheries production: A review. *Archives of Applied Science Research* 8(6): 44–51.
- Paul B N, Chanda S, Bhowmick S, Sridhar N, Saha G S and Giri S S. (2017). Nutrient profile of Indian climbing perch, *Anabas testudineus*. *SAARC Journal of Agriculture* 15(1): 99–109. <https://doi.org/10.3329/sja.v15i1.33156>
- Pauly D. (2019). Female fish grow bigger – Let’s deal with it. *Trends in Ecology & Evolution TREE* 34(3): 181–182.
- Perez-Palafox X A, Morales-Bojórquez E, Aguirre-Villaseñor H and Cruz-Escalona V H. (2022). Length at maturity, sex ratio, and proportions of maturity of the giant electric ray, *Narcine entemedor*, in its septentrional distribution. *Animals* 12(1): 120. <https://doi.org/10.3390/ani12010120>
- Perlman H. (2013). Water properties: Temperature. In the USGS Water Science School. <https://ga.water.usgs.gov/edu/temperature.html> (Accessed on 1 April 2019)
- Petersen C W and Warner R R. (2002). The ecological context of reproductive behavior. In P F Sale (ed.), *Coral reef fishes: Dynamics and diversity in a complex ecosystem*. San Diego: Academic Press, 103–122. <https://doi.org/10.1016/B978-012615185-5/50007-4>
- Piferrer F and Guiguen Y. (2008). Fish gonadogenesis. Part II: Molecular biology and genomics of sex differentiation. *Reviews in Fisheries Science* 16(sup. 1): 35–55. <https://doi.org/10.1080/10641260802324644>
- Priyatha C V and Chitra K C. (2022). Evaluation of the reproductive cycle and gonadal development in the climbing perch, *Anabas testudineus* (Bloch, 1792) in captivity. *Journal of Fisheries* 10(1): 101206–101206. <https://doi.org/10.17017/j.fish.364>
- Rahman M A and Marimuthu K. (2010). Effect of different stocking density on growth, survival and production of endangered native fish climbing perch (*Anabas testudineus*, Bloch) fingerlings in nursery ponds. *Advances in Environmental Biology* 4(2): 178–187.
- Rahman M M and Samat A F. (2021). Reproductive cycle, sexual maturity and fecundity of *Nemipterus furcosus* (Valenciennes, 1830). *Aquaculture and Fisheries* 6(4): 424–431. <https://doi.org/10.1016/j.aaf.2020.07.006>
- Rathod S R, Shinde S E and More S R. (2011). Length-weight relationship in *Salmostoma navacula* and *Channa maurulius* Godavari River at Kaigaon, Toka, Dist. Aurangabad (M.S.) India. *Recent Research in Science and Technology* 3(3): 104–106.
- Russell M W, Luckhurst B E and Lindeman K C. (2011). Management of spawning aggregations. In Y S de Mitcheson and P L Colin (eds.), *Reef fish spawning aggregations: biology, research and management*. Dordrecht: Springer Netherlands, 371–404. https://doi.org/10.1007/978-94-007-1980-4_11

- Sarkar U K, Deepak P K, Kapoor D, Negi R S, Paul S K and Singh S. (2005). Captive breeding of climbing perch, *Anabas testudineus* (Bloch, 1792) with Wova-FH for conservation and aquaculture. *Aquaculture Research* 36(10): 941–945. <https://doi.org/10.1111/j.1365-2109.2005.01281.x>
- Schuler B. (2014). *Environmental and climate change in South and Southeast Asia: How are local cultures coping?* Leiden, The Netherlands: Koninklijke Brill NV Publisher. <https://doi.org/10.1163/9789004273221>
- Schulz R W, Renato D F, Luiz, Jean-Jacques L, Florence L, Chiarini-Garcia H, Nobrega R H and Miura T. (2010). Spermatogenesis in fish. *General and Comparative Endocrinology* 165: 390–411. <https://doi.org/10.1016/j.ygcen.2009.02.013>
- Shahriar F, Ilnasina A A, Uddin N, Munny F J, Hussain M A, Jang W J, Lee J M, Lee E W, Hasan M T and Kawsar M A. (2024). Effects of commercial probiotics on the growth, hematology, immunity, and resistance to *Aeromonas hydrophila* challenge in climbing perch, *Anabas testudineus*. *Aquaculture Research* 1: 3901035. <https://doi.org/10.1155/are/3901035>
- Stasko A D, Swanson H, Majewski A, Atchison S, Reist J and Power M. (2016). Influences of depth and pelagic subsidies on the size-based trophic structure of Beaufort Sea fish communities. *Marine Ecology Progress Series* 549: 153–166. <https://doi.org/10.3354/meps11709>
- Suchiang P and Gupta B B P. (2011). Variations in the plasma levels of thyroid hormones and testicular activity in the male air-breathing catfish (*Clarias gariepinus*) over the annual cycle. *International Journal of Biological Sciences* 3(3): 144–154. <https://doi.org/10.5539/ijb.v3n3p144>
- Sumon M S, Ali M R and Sarower M G. (2015). The effects of water quality additives on bacterial load and production in Thai climbing perch (*Anabas testudineus* Bloch, 1792) culture pond. *International Journal of Fisheries and Aquatic Studies* 2(6): 235–238.
- Syazwani N. (2020). RM20 per kilogram of hybrid climbing perch; High potential livestock [online article]. *Rentas Asia*. <https://rentas.asia/rm20-sekilogram-ikan-puyu-hibrid-ternakan-yang-berpotensi-tinggi/>
- Tay Y L, Loong A M, Hiong K C, Lee S J J, Tng Y Y M, Wee N L J, Lee S M L, Wang W P, Chew S F, Wilson J M and Ip Y K. (2006). Active ammonia transport and excretory nitrogen metabolism in the climbing perch, *Anabas testudineus*, during 4 days of emersion or 10 minutes of forced exercise on land. *Journal of Experimental Biology* 209: 4475–4489. <https://doi.org/10.1242/jeb.02557>
- Thompson L and Larsen R. (2004). *Fish habitat in freshwater streams*. UCANR Publications. <https://doi.org/10.3733/ucanr.8112>
- Tian P, Yang Z, Qu C, Qi X, Zhu L, Hao G and Zhang Y (2024). Exploration of tissue fixation methods suitable for digital pathological studies of the testis. *European Journal of Medical Research* 29(1): 319. <https://doi.org/10.1186/s40001-024-01921-5>
- Tran K C, Van Thuan D and Nguyen T L. (2024). Antimicrobial activity of some plant extracts against *Kosakonia sacchari* causing diseases on a Vietnamese climbing perch (*Anabas testudineus*). *IOP Conference Series: Earth and Environmental Science* 1340(1): 012019. <https://doi.org/10.1088/1755-1315/1340/1/012019>
- Uddin K B, Moniruzzaman M, Bashar M A, Basak S, Islam A S, Mahmud Y, Lee S and Bai S C. (2016). Culture potential of Thai climbing perch (*Anabas testudineus*) in experimental cages at different stocking densities in Kaptai Lake, Bangladesh. *Aquaculture, Aquarium, Conservation & Legislation* 9(3): 564–573.

- Ultsch G R. (1989). Ecology and physiology of hibernation and overwintering among freshwater fishes, turtle and snakes. *Biological Reviews* 64: 435–515. <https://doi.org/10.1111/j.1469-185X.1989.tb00683.x>
- Uttam K.S, Prashant K D, Dhurenda K, Raje S N, Samir K P and Sreepakash S. (2005). Captive breeding of climbing perch, *Anabas testudineus* with Wova-FH for conservation and aquaculture. *Aquaculture Research* 36: 941–945. <https://doi.org/10.1111/j.1365-2109.2005.01281.x>
- Wilson S K, Fisher R, Pratchett M S, Graham N A J, Dulvy N K, Turner R A, Cakacaka A and Polunin N V. (2010). Habitat degradation and fishing effects on the size structure of coral reef fish communities. *Ecological Applications* 20(2): 442–451. <https://doi.org/10.1890/08-2205.1>
- Yoneda M and Wright P J. (2005). Effect of temperature and food availability on reproductive investment of first-time spawning male Atlantic cod, *Gadus morhua*. *ICES Journal of Marine Science* 62(7): 1387–1393. <https://doi.org/10.1016/j.icesjms.2005.04.018>
- Zalina I, Saad, C R, Christianus A and Harmin S A. (2012). Induced breeding and embryonic development of climbing perch (*Anabas testudineus*, Bloch). *Journal of Fisheries and Aquatic Science* 7: 291–306. <https://doi.org/10.3923/jfas.2012.291.306>
- Zworykin D D. (2012). Reproduction and spawning behavior of climbing perch, *Anabas testudineus* (Perciformes, Anabantidae) in an Aquarium. *Journal of Ichthyology* 52(6): 379–388. <https://doi.org/10.1134/S0032945212040169>